

Forestation in hypergraphs: linear k -trees

Ojas Parekh

Department of Mathematics and Computer Science
Emory University, Atlanta, USA
ojas@mathcs.emory.edu

Submitted: September 14, 2001; Accepted: September 1, 2003.

MR Subject Classifications: 05C65, 05E99

Abstract

We present a new proof of a result of Lovász on the maximum number of edges in a k -forest. We also apply a construction used in our proof to generalize the notions of a k -hypertree and k -forest to a class which extends some properties of trees, to which both specialize when $k=2$.

1 Introduction

Let $X = [n]$ and \mathcal{F} be a k -uniform hypergraph on X . We say an edge $e \in \mathcal{F}$ *crosses* a k -partition, $X = X_1 \dot{\cup} \dots \dot{\cup} X_k$, if $|e \cap X_i| = 1$ for $1 \leq i \leq k$. \mathcal{F} is a k -forest if for each $e \in \mathcal{F}$ there is some k -partition $X = X_1^e \dot{\cup} \dots \dot{\cup} X_k^e$ such that e is the unique edge crossing it. What is the maximum number of edges in \mathcal{F} ?

This problem was initially posed to László Lovász by Ronald Graham [2]. Lovász's novel algebraic proof appeared in [3] in 1979, and our proof remains algebraic in nature; however, it relies on homogeneous multilinear polynomials over \mathbb{F}_2 rather than tensors. The reader is encouraged to consult [1] for an introduction to and extensive applications of linear algebra in combinatorics.

Theorem 1.1. *A k -forest \mathcal{F} on X has at most $\binom{n-1}{k-1}$ edges.*

Proof. We open with a few definitions. By \mathbb{P}_{k-1}^{n-1} we mean the space of multilinear homogeneous polynomials of degree $k-1$ in $\mathbb{F}_2[x_1, \dots, x_{n-1}]$. We make use of the shorthand $p(\mathbf{x})$ to denote $p(x_1, \dots, x_{n-1})$, where $\mathbf{x} = (x_1, \dots, x_{n-1}) \in \mathbb{F}_2^{n-1}$ and $p \in \mathbb{P}_{k-1}^{n-1}$. Finally, for $e \in \mathcal{F}$, $\mathbb{1}_e$ denotes the incidence vector of e .

For each edge $e \in \mathcal{F}$ we pick a k -partition $\pi_e = (X_1^e, \dots, X_k^e)$, such that e is the unique edge crossing it. For simplicity we assume X_1^e contains the element n . We then define a polynomial,

$$p_e(x_1, \dots, x_{n-1}) = \prod_{i=2}^k \sum_{j \in X_i^e} x_j.$$

For each e in \mathcal{F} , p_e is in \mathbb{P}_{k-1}^{n-1} , hence it suffices to demonstrate the independence of these polynomials. To that end we seek to show that if $e, f \in \mathcal{F}$, then $p_e(\mathbb{1}_{f \setminus \{n\}}) = 1$ if and only if $f = e$. We have

$$p_e(\mathbb{1}_{f \setminus \{n\}}) = \prod_{i=2}^k (|f \cap X_i^e| \bmod 2).$$

Clearly $p_e(\mathbb{1}_{e \setminus \{n\}}) = 1$. If $f \neq e$ there must be some i for which $|f \cap X_i^e| = 0$, since f does not cross π_e . In this case there also exists a $j \neq i$ such that $|f \cap X_j^e| \bmod 2 = 0$. Thus $p_e(\mathbb{1}_{f \setminus \{n\}}) = 0$. \square

Our agenda for the remainder of the paper is to first consider a generalization of k -forests which preserves certain properties of forests and to then proceed to compare our generalization with existing ones.

2 Linear k -trees

In light of the result of the previous section a natural question arises. What can one say about the maximum k -forests, those with exactly $\binom{n-1}{k-1}$ edges? We could begin by considering small examples. It is not difficult to verify that a 2-forest is indeed a forest. In this case any maximal forest is a tree, which one may define in several ways. A basic result in graph theory is that a graph which exhibits any two of

- (i) acyclicity
- (ii) exactly $n - 1$ edges
- (iii) connectivity

necessarily exhibits the third.

We already have analogues of (i) and (ii) that we could use in defining a k -tree for $k > 2$, and one might conjecture that for a k -uniform hypergraph \mathcal{H} on X , any two of

- (i') \mathcal{H} is a k -forest.
- (ii') \mathcal{H} has exactly $\binom{n-1}{k-1}$ edges.
- (iii') For each k -partition of X , \mathcal{H} contains an edge that crosses it.

implies the third. Unfortunately this is not true.

Counterexample 2.1. *The 3-uniform hypergraph*

$$\mathcal{H} = \{\{1, 2, 4\}, \{1, 2, 5\}, \{1, 3, 5\}, \{1, 4, 5\}, \{2, 3, 4\}, \{3, 4, 5\}\}$$

over $\{1, 2, 3, 4, 5\}$ satisfies (ii') and (iii') but not (i').

Why does this generalization fail? Conditions (i')-(iii') extend the notion of a cut in a graph, which is implicit in (i)-(iii), to that of a k -partition. We do have that a 2-partition is indeed a cut; however, this is not the whole story. The proof of Theorem 1.1 offers some insight into the matter. The multilinear polynomial space \mathbb{P}_1^{n-1} consists entirely of polynomials which correspond to 2-partitions; however, the reader may verify that an analogous statement is not true for even \mathbb{P}_2^{n-1} . Guided by this discrepancy, we say an edge $e \in \mathcal{H}$ *crosses* a *polynomial* $p \in \mathbb{P}_{k-1}^{n-1}$ if $p(\mathbb{1}_{e \setminus \{n\}}) = 1$, and we relax (i'): The hypergraph \mathcal{H} is a *linear k -forest* (vs. a k -forest) if for each edge $e \in \mathcal{H}$, there is a polynomial $p_e \in \mathbb{P}_{k-1}^{n-1}$ (vs. a k -partition) such that e is the unique edge in \mathcal{H} crossing p_e . We accordingly strengthen (iii'): The hypergraph \mathcal{H} is *linearly k -connected*, or simply *k -connected*, if for each polynomial $p \in \mathbb{P}_{k-1}^{n-1}$, there is an edge $e \in \mathcal{H}$ which crosses p . The scrutinizing reader might have sensed something amiss in the preceding definitions. The polynomial space \mathbb{P}_{k-1}^{n-1} is defined with respect to a distinguished element $n \in X$.

Lemma 2.2. *A hypergraph is a linear k -forest or k -connected independently of the choice of distinguished element used in defining \mathbb{P}_{k-1}^{n-1} .*

Proof. Let $p(x_1, \dots, x_{n-1}) \in \mathbb{P}_{k-1}^{n-1}$. We will demonstrate a $p'(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n) \in \mathbb{P}_{k-1}^n$ such that $\{e \in \binom{X}{k} \mid p(\mathbb{1}_{e \setminus \{n\}}) = 1\} = \{e \in \binom{X}{k} \mid p'(\mathbb{1}_{e \setminus \{i\}}) = 1\}$. We divide p by x_i to yield $p = x_i q + r$ where $q \in \mathbb{P}_{k-2}^{n-1}$, $r \in \mathbb{P}_{k-1}^{n-1}$, and neither contain the variable x_i . We can represent q as a sum of monomials, that is there exist sets $Y_s \in \binom{X \setminus \{i, n\}}{k-2}$ for s in some index set S such that $q = \sum_{s \in S} \prod_{j \in Y_s} x_j$. Notice that an edge crosses the polynomial $(\prod_{j \in Y_s} x_j)(\sum_{j \notin Y_s \cup \{i\}} x_j)$ if and only if it crosses the monomial $x_i(\prod_{j \in Y_s} x_j)$. This provides us the construction we seek, and we set

$$p'(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n) = r + \sum_{s \in S} \left(\prod_{j \in Y_s} x_j \right) \left(\sum_{j \notin Y_s \cup \{i\}} x_j \right).$$

□

We will henceforth use \mathbb{P}_{k-1}^{n-1} to refer to a multilinear polynomial space in $n-1$ variables, the indices of which will be clear from context. We now have the following.

Theorem 2.3. *For \mathcal{H} , a k -uniform hypergraph on X , any two of*

- (i) \mathcal{H} is a linear k -forest.
- (ii) \mathcal{H} has exactly $\binom{n-1}{k-1}$ edges.
- (iii) \mathcal{H} is k -connected.

implies the third.

Proof.

(i),(ii) implies (iii): For each edge e let p_e be a polynomial for which e is the unique edge in \mathcal{H} crossing p_e . For a polynomial $p \in \mathbb{P}_{k-1}^{n-1}$, let $\mathcal{H}(p)$ denote $\{e \in \mathcal{H} \mid e \text{ crosses } p\}$. As in the proof of Theorem 1.1 we have the independence of the polynomials p_e for $e \in \mathcal{H}$, hence $|\{p_e \mid e \in \mathcal{H}\}| = \binom{n-1}{k-1}$ by (ii). The set $\{p_e \mid e \in \mathcal{H}\}$ is a basis for \mathbb{P}_{k-1}^{n-1} , so for any $q \in \mathbb{P}_{k-1}^{n-1}$ we must have $\mathcal{H}(q) \neq \emptyset$.

(ii),(iii) implies (i): First we establish $p \neq q$ implies $\mathcal{H}(p) \neq \mathcal{H}(q)$, for polynomials $p, q \in \mathbb{P}_{k-1}^{n-1}$. Proceeding by contrapositive, if $\mathcal{H}(p) = \mathcal{H}(q)$ then $\mathcal{H}(p+q) = \emptyset$, hence $p = q$. There are exactly $2^{\binom{n-1}{k-1}} - 1$ polynomials in \mathbb{P}_{k-1}^{n-1} and $|\mathcal{H}| = \binom{n-1}{k-1}$, so by (iii) $\{\mathcal{H}(p) \mid p \in \mathbb{P}_{k-1}^{n-1}\} = 2^{\mathcal{H}} \setminus \{\emptyset\}$, where $2^{\mathcal{H}}$ is the powerset of \mathcal{H} . Thus for each edge $e \in \mathcal{H}$, $\{e\} \in \{\mathcal{H}(p) \mid p \in \mathbb{P}_{k-1}^{n-1}\}$.

(iii),(i) implies (ii): From the proof of the first part we have (i) implies $|\mathcal{H}| \leq \binom{n-1}{k-1}$; from that of the second we have (iii) implies $|\mathcal{H}| \geq \binom{n-1}{k-1}$.

□

We are finally in position to call a hypergraph \mathcal{T} that satisfies any two conditions above a linear k -tree. The third part of the proof of the theorem hints at two other characterizations of linear k -trees.

Theorem 2.4.

- (i) *Every k -connected hypergraph contains a linear k -tree.*
- (ii) *Every linear k -forest is contained in a linear k -tree.*

Proof.

(i): For the sake of contradiction, let \mathcal{H} be a minimal k -uniform hypergraph over X that is k -connected but does not contain a linear k -tree. We let $\mathbb{P}(H)$ represent $\{p \in \mathbb{P}_{k-1}^{n-1} \mid \text{some } e \in H \text{ crosses } p\}$, where $H \subseteq \mathcal{H}$; we omit braces for singleton arguments. Since \mathcal{H} is not a linear k -forest, there is some $e \in \mathcal{H}$ such that $\mathbb{P}(e) \subseteq \mathbb{P}(\mathcal{H} \setminus \{e\})$, hence $\mathcal{H} \setminus \{e\}$ is also a counterexample.

(ii): For the sake of contradiction, let \mathcal{H} be a maximal k -uniform hypergraph over X that is a linear k -forest but is not contained in a linear k -tree. Since \mathcal{H} is not k -connected, there is some $p \in \mathbb{P}_{k-1}^{n-1}$ such that $\mathcal{H}(p) = \emptyset$. Let $f \in \binom{X}{k}$ be some set such that $p(\mathbb{1}_{f \setminus \{n\}}) = 1$, and for $e \in \mathcal{H}$ let p_e be a polynomial such that $\mathcal{H}(p_e) = \{e\}$. We set $p'_f = p$, and for each edge $e \in \mathcal{H}$, we set

$$p'_e = \begin{cases} p_e + p & \text{if } p_e(\mathbb{1}_{f \setminus \{n\}}) = 1 \\ p_e & \text{otherwise} \end{cases},$$

which renders e the unique edge in $\mathcal{H} \cup \{f\}$ crossing p'_e and $\mathcal{H} \cup \{f\}$ a counterexample.

□

Thus we may also think of linear k -trees as maximal linear k -forests or minimally k -connected hypergraphs.

3 All trees are not created equal

A linear k -tree is only one of a multitude of possible generalizations of trees to hypergraphs; in this section we explore the connection between linear k -trees and a generalization which exists in the literature.

The combinatorial structure known as a k -hypertree was introduced in [4] as a tool for developing Bonferroni type inequalities. A k -hypertree is a k -uniform hypergraph \mathcal{T} on X such that for $k = 2$, \mathcal{T} is a tree with vertex set X and for $k \geq 3$, \mathcal{T} is defined recursively as follows:

- (i) If $X = \{1, \dots, k\}$ then \mathcal{T} has a unique edge $\{1, \dots, k\}$.
- (ii) If $|X| \geq k + 1$ then there exists an element $i \in X$ such that if e_1, \dots, e_q denote all edges containing i then $e_1 \setminus \{i\}, \dots, e_q \setminus \{i\}$ induce an $(k - 1)$ -hypertree with vertex set $X \setminus \{i\}$ and the remaining edges of \mathcal{T} induce a k -hypertree with vertex set $X \setminus \{i\}$. A k -hypertree has exactly $\binom{n-1}{k-1}$ edges.

The notion was augmented [5] by imposing a total ordering μ on X , yielding several nice characterizations of k -hypertrees which generalize properties of trees. We show that linear k -trees generalize k -hypertrees. We denote the classes of linear k -trees and k -hypertrees on X by $\mathcal{LKT}(k, n)$ and $\mathcal{HT}(k, n)$ respectively.

Theorem 3.1. $\mathcal{HT}(k, n) \subset \mathcal{LKT}(k, n)$.

Proof. We show inclusion by induction. We have that $\mathcal{HT}(k, k) = \mathcal{LKT}(k, k)$, so let us consider some $\mathcal{T} \in \mathcal{HT}(k, n)$ for $k < n$. Since $|\mathcal{T}| = \binom{n-1}{k-1}$, by Theorem 2.3 it suffices to show \mathcal{T} is k -connected. Let $l \in X$ be an element such that $\mathcal{T}_l = \{e \setminus \{l\} \mid l \in e \in \mathcal{T}\}$ and $\mathcal{T}_{\bar{l}} = \{e \in \mathcal{T} \mid l \notin e\}$ are respectively $(k - 1)$ - and k -hypertrees over $X \setminus \{l\}$.

We seek to show that for a polynomial $p(x_1, \dots, x_{n-1}) \in \mathbb{P}_{k-1}^{n-1}$ there is some edge in $\mathcal{T} = \mathcal{T}_l \cup \mathcal{T}_{\bar{l}}$ that crosses it. Note that we may assume $l \neq n$ by Lemma 2.2. Dividing by x_l gives us $p = x_l q(x_1, \dots, x_{l-1}, x_{l+1}, \dots, x_{n-1}) + r(x_1, \dots, x_{l-1}, x_{l+1}, \dots, x_{n-1})$. If $r \equiv 0$ then $e \cup \{l\}$ crosses $p = x_l q$ for some $e \in \mathcal{T}_l$, since $\mathcal{T}_l \in \mathcal{KT}(k - 1, n - 1)$ by the induction hypothesis. Otherwise some $e \in \mathcal{T}_{\bar{l}}$ crosses r , since $\mathcal{T}_{\bar{l}} \in \mathcal{KT}(k, n - 1)$ by the induction hypothesis. In this case $l \notin e$, hence e crosses $p = x_l q + r$.

As for strict inclusion, we leave it to the reader to verify that

$$\mathcal{T} = \left\{ \{1, 2, 3\}, \{1, 2, 4\}, \{1, 2, 6\}, \{1, 4, 5\}, \{1, 5, 6\}, \right. \\ \left. \{2, 3, 5\}, \{2, 3, 6\}, \{3, 4, 5\}, \{3, 4, 6\}, \{4, 5, 6\} \right\}$$

is a linear 3-tree but not a 3-hypertree. □

The class $\mathcal{LKT}(k, n)$ may be a practically significant generalization of $\mathcal{HT}(k, n)$. Given a cost function $c : \binom{X}{k} \rightarrow \mathbb{R}_+$, it is NP -complete to decide whether there is a k -hypertree of cost at most l for $n > k \geq 3$ [6]. This is known as the minimum spanning k -hypertree

problem and for $k = 2$ reduces to the polynomial time solvable minimum spanning tree problem. Replacing ‘ k -hypertree’ with ‘linear k -tree’ in the above definition drastically reduces the complexity of the problem. By Theorems 2.4 and 2.3 the linear k -forests on X comprise a matroid, hence we can apply a greedy algorithm to solve the minimum spanning linear k -tree problem in polynomial time for constant k .

We close by offering a conjecture. A k -tree is a k -forest of size $\binom{n-1}{k-1}$. We let $\mathcal{KT}(k, n)$ denote the class of k -trees on X . From Theorem 2.3, Counterexample 2.1, and the fact that $\{e \in \binom{X}{k} \mid 1 \in e\} \in \mathcal{HT}(k, n) \cap \mathcal{KT}(k, n)$, we derive the following properties.

$$\mathcal{KT}(k, n) \subset \mathcal{LKT}(k, n) \tag{1}$$

$$\mathcal{HT}(k, n) \setminus \mathcal{KT}(k, n) \neq \emptyset \tag{2}$$

$$\mathcal{HT}(k, n) \cap \mathcal{KT}(k, n) \neq \emptyset \tag{3}$$

Unfortunately these leave the precise interaction of $\mathcal{HT}(k, n)$ and $\mathcal{KT}(k, n)$ uncertain. Yet if one could show that for every $\mathcal{T} \in \mathcal{KT}(k, n)$ there is some $i \in X$ that is contained in exactly $\binom{n-2}{k-2}$ edges, then induction would yield the following.

Conjecture 3.2. $\mathcal{KT}(k, n) \subset \mathcal{HT}(k, n)$.

Acknowledgements

The author is grateful to Tom Bohman for exposing him to the tools and the trade.

References

- [1] L. Babai and P. Frankl. *Linear Algebra Methods in Combinatorics*. Dept. of Computer Science, The Univ. of Chicago, Chicago, 1992.
- [2] R. Graham. Personal Communication. 2000.
- [3] L. Lovász. Topological and Algebraic Methods in Graph Theory. In *Graph theory and related topics (Proc. Conf., Univ. Waterloo, Waterloo, Ont., 1977)*, pp. 1–14, Academic Press, New York-London, 1979.
- [4] I. Tomescu. Hypertrees and Bonferroni inequalities. *J. Combin. Theory Ser. B*, 41:209–217, 1986.
- [5] I. Tomescu. Ordered h-Hypertrees. *Discrete Mathematics*, 195:241–248, 1992.
- [6] I. Tomescu and M. Zimand. Minimum spanning hypertrees. *Discrete Appl. Math.*, 54.1:67–76, 1994.