

IDENTITIES FOR TRACES OF SINGULAR MODULI

KATHRIN BRINGMANN AND KEN ONO

ABSTRACT. Generalizing work of Zagier, in an important recent paper Bruinier and Funke prove that the generating functions for traces of singular values of many modular functions are weight $\frac{3}{2}$ modular forms. Using facts about half-integral weight modular forms, we obtain identities relating traces of singular moduli for modular functions of level p and 1. These follow from a general result relating coefficients of certain weakly holomorphic weight $\frac{3}{2}$ modular forms of level $4p$ to Zagier's traces of singular moduli.

1. INTRODUCTION AND STATEMENT OF RESULTS

Let $q := e^{2\pi iz}$ and let

$$J(z) = q^{-1} + 196884q + 21493760q^2 + \dots$$

be the Hauptmodul for $\Gamma = \mathrm{PSL}_2(\mathbb{Z})$. Similarly, let

$$j_3^*(z) = q^{-1} + 783q + 8672q^2 + 65367q^3 + 371520q^4 + \dots$$

be the Hauptmodul for $\Gamma_0^*(3)$, the extension of $\Gamma_0(3)$ by the Fricke involution W_3 . The values of such functions at imaginary quadratic arguments are known as *singular moduli*.

There are general identities relating singular moduli for functions such as $J(z)$ and $j_3^*(z)$. To illustrate, observe that the evaluations $J\left(\frac{-1+\sqrt{-3}}{2}\right) = -744$, $j_3^*\left(\frac{-3+\sqrt{-3}}{6}\right) = -42$, and $j_3^*\left(\frac{-7+\sqrt{-3}}{6}\right) = -234$ imply that

$$(1.1) \quad J\left(\frac{-1+\sqrt{-3}}{2}\right) = j_3^*\left(\frac{-3+\sqrt{-3}}{6}\right) + 3j_3^*\left(\frac{-7+\sqrt{-3}}{6}\right).$$

Such identities follow from arithmetic relations between modular forms of different levels.

For a positive integer m , let $J_m(z)$ be the unique modular function on Γ with Fourier expansion

$$J_m(z) = q^{-m} + \sum_{n=1}^{\infty} c_m(n)q^n.$$

Zagier [Za] computed the generating functions for the “traces” of the $J_m(z)$ singular moduli, as well as several other classes of modular functions. To describe these traces, let

Date: June 6, 2005.

2000 Mathematics Subject Classification. 11F37, 11F30.

The second author thanks the Packard Foundation, the Guggenheim Foundation, the support of a Romnes Fellowship, and the National Science Foundation.

$D \equiv 0, 3 \pmod{4}$ be a positive integer, and let \mathcal{Q}_D denote the positive definite integral binary quadratic forms $Q(a, b, c) = ax^2 + bxy + cy^2$ of discriminant $-D = b^2 - 4ac$. The group Γ acts on \mathcal{Q}_D by

$$Q \circ \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} := Q(\alpha x + \beta y, \gamma x + \delta y).$$

For each $Q \in \mathcal{Q}_D$, we let $\alpha_Q := \frac{-b + \sqrt{b^2 - 4ac}}{2a} \in \mathbb{H}$, and we let $\omega_Q := |\Gamma_Q|$, where Γ_Q is the isotropy subgroup of Q in Γ . If m is a positive integer and $-D$ is a discriminant, then Zagier defined the trace of the singular moduli of discriminant $-D$ for $J_m(z)$ by

$$(1.2) \quad t_m(D) := \sum_{Q \in \mathcal{Q}_D/\Gamma} \frac{1}{\omega_Q} \cdot J_m(\alpha_Q).$$

He proved the striking fact that their generating functions are weight $\frac{3}{2}$ modular forms.

It turns out that (1.1) is a special case of a general phenomenon where sums of coefficients of certain weakly holomorphic weight $\frac{3}{2}$ modular forms are given in terms of such traces. A meromorphic modular form is *weakly holomorphic* if it is holomorphic on \mathbb{H} . Following Kohnen, for integers k let $\mathcal{M}_{k+\frac{1}{2}}^+(\Gamma_0(4))$ be the space of weakly holomorphic weight $k + \frac{1}{2}$ modular forms on $\Gamma_0(4)$ with a Fourier expansion of the form

$$(1.3) \quad \sum_{\substack{n \gg -\infty \\ (-1)^k n \equiv 0, 1 \pmod{4}}} a(n)q^n.$$

Furthermore, if p is an odd prime and $\epsilon = \pm 1$, then let $\mathcal{M}_{k+\frac{1}{2}}^{+, \epsilon}(\Gamma_0(4p))$ be the space of those weight $k + \frac{1}{2}$ weakly holomorphic modular forms $f(z) = \sum_{n \gg -\infty} a(n)q^n$ on $\Gamma_0(4p)$ whose Fourier coefficients satisfy

$$(1.4) \quad a(n) = 0 \quad \text{whenever} \quad (-1)^k n \equiv 2, 3 \pmod{4} \quad \text{or} \quad \left(\frac{(-1)^k n}{p} \right) = -\epsilon.$$

The generic identities, for forms in $\mathcal{M}_{\frac{3}{2}}^{+, \epsilon}(\Gamma_0(4p))$, are expressed in terms of a distinguished sequence of modular forms. To define them, for each positive $m \equiv 0, 1 \pmod{4}$ let $g_m(z)$ be the unique form in $\mathcal{M}_{\frac{3}{2}}^+(\Gamma_0(4))$ with a Fourier expansion of the form

$$(1.5) \quad g_m(z) = q^{-m} + \sum_{n=0}^{\infty} B(m, n)q^n.$$

Remark. The uniqueness of the $g_m(z)$ essentially follows from the fact that there are no holomorphic forms in $\mathcal{M}_{\frac{3}{2}}^+(\Gamma_0(4))$ (see the discussion preceding Theorem 4 of [Za]).

Theorem 1.1. *If p is an odd prime, $\epsilon = \pm 1$, and $g = \sum_{n \gg -\infty} a(n)q^n \in \mathcal{M}_{\frac{3}{2}}^{+, \epsilon}(\Gamma_0(4p))$, then for every positive $n \equiv 0, 3 \pmod{4}$ we have*

$$a(n) + \epsilon a(np^2) = \sum_{m \geq 1} (a(-m) + \epsilon a(-mp^2)) B(m, n).$$

Remark. Since Zagier (see Section 7 of [Za]) proved that the integers $B(m, n)$ are traces or “twisted traces” of singular moduli, Theorem 1.1 provides identities expressing sums of coefficients of forms in $\mathcal{M}_{\frac{3}{2}}^{+, \epsilon}(\Gamma_0(4))$ as sums of traces of singular moduli. For brevity and aesthetics, we do not give the general formulas here. Instead, we simply give Corollary 1.2 which relates the traces $t_m(D)$ to the level p traces defined below.

Recently, Bruinier and Funke [BF] have greatly generalized Zagier’s results to include traces of singular moduli of modular functions on groups which do not necessarily possess a Hauptmodul. A particularly elegant example of their general work corresponds to modular functions on $\Gamma_0(p)$ which are fixed by the Fricke involution $W_p = \begin{pmatrix} 0 & -1 \\ p & 0 \end{pmatrix}$. To define these traces, let p be an odd prime, and let $\Gamma_0^*(p)$ be the projective image of the extension of $\Gamma_0(p)$ by the Fricke involution W_p in $\mathrm{PSL}_2(\mathbb{R})$. Let $\mathcal{Q}_{D,p}$ be the set of quadratic forms $Q \in \mathcal{Q}_D$ such that $a \equiv 0 \pmod{p}$. The group $\Gamma_0^*(p)$ acts on $\mathcal{Q}_{D,p}$, where the action for elements of $\Gamma_0(p)$ is defined as before, and $Q \circ W_p := Q'(x, y)$, where $Q'(x, y) = (pc, -b, a/p)$.

Suppose that $f(z)$ is in $\mathcal{M}_0(\Gamma_0^*(p))$, the set of weakly holomorphic modular functions for $\Gamma_0^*(p)$. For such f , the discriminant $-D$ trace is given by

$$(1.6) \quad t_f^*(D) := \sum_{Q \in \mathcal{Q}_{D,p}/\Gamma_0^*(p)} \frac{1}{|\Gamma_0^*(p)_Q|} \cdot f(\alpha_Q).$$

Here $\Gamma_0^*(p)_Q$ is the stabilizer of Q in $\Gamma_0^*(p)$. Applying Theorem 1.1 to the Bruinier-Funke generating functions gives the following corollary.

Corollary 1.2. *If p is an odd prime and $f = \sum a(n)q^n \in \mathcal{M}_0(\Gamma_0^*(p))$, with $a(0) = 0$, then for every positive integer $D \equiv 0, 3 \pmod{4}$ we have*

$$t_f^*(D) + t_f^*(Dp^2) = \sum_{m \geq 1} \frac{b(m)}{m} \sum_{l|m} \mu(m/l) t_l(D).$$

Here $\mu(n)$ denotes the usual Möbius function, and

$$b(m) = \sum_{n \geq 1} (ma(-mn) + p^2 ma(-p^2 mn)).$$

Remark. There is an alternate approach to proving Corollary 1.2. Using the weight zero analog of Lemma 2.2, one can obtain an identity of the form

$$\sum_{m=1}^M c(m) J_m(z) = f(z) + pf(z)|U(p).$$

Using this identity, one easily obtains sums of $t_m(n)$ as sums of singular moduli for $f(z)$. To prove Corollary 1.2, one needs to relate such sums to (1.6). This requires a somewhat lengthy calculation which aligns the corresponding CM points, counted with the correct multiplicity, with the points obtained from definition of the $U(p)$ operator. The extra information provided by Bruinier and Funke, and Zagier, allows us to argue instead with standard properties of half-integral weight modular forms.

Example. If $f = j_3^*$, $D = 3$, and $p = 3$, then Corollary 1.2 implies that

$$t_{j_3^*}^*(3) + t_{j_3^*}^*(27) = t_1(3) = \frac{1}{3} \cdot J \left(\frac{-1 + \sqrt{-3}}{2} \right) = -248.$$

It turns out that $t_{j_3^*}^*(3) = \frac{j_3^* \left(\frac{-3 + \sqrt{-3}}{6} \right)}{6} = -\frac{42}{6} = -7$ and

$$t_{j_3^*}^*(27) = j_3^* \left(\frac{-7 + \sqrt{-3}}{6} \right) + \frac{j_3^* \left(\frac{-9 + \sqrt{-3}}{6} \right)}{6} = -234 - \frac{42}{6} = -241.$$

Noting that $j_3^* \left(\frac{-3 + \sqrt{-3}}{6} \right) = j_3^* \left(\frac{-9 + \sqrt{-3}}{6} \right) = -42$ gives (1.1).

In Section 2 we recall facts about half-integral weight modular forms. In Section 3 we discuss features of the ‘‘Kohnen spaces’’, recall the formulation of the Bruinier-Funke and the Zagier generating functions, and prove Theorem 1.1 and Corollary 1.2.

ACKNOWLEDGEMENTS

The authors thank W. Kohnen and J. Rouse for their helpful comments.

2. PRELIMINARIES ON HALF-INTEGRAL WEIGHT MODULAR FORMS

We recall basic facts about half-integral weight modular forms (see [Sh] and [Ko]). Throughout, let k be an integer, and let $p = 1$ or an odd prime. We let $\mathfrak{S}_{k+\frac{1}{2}}$ be the group consisting of all pairs $(A, \phi(z))$, where $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{GL}_2^+(\mathbb{R})$ and ϕ is a complex-valued holomorphic function on \mathbb{H} satisfying

$$|\phi(z)| = (\det A)^{-\frac{k}{2} - \frac{1}{4}} |cz + d|^{k+\frac{1}{2}}.$$

The group law in $\mathfrak{S}_{k+\frac{1}{2}}$ is given by

$$(2.1) \quad (A, \phi(z)) \cdot (B, \psi(z)) := (AB, \phi(Bz)\psi(z)).$$

The group algebra of $\mathfrak{S}_{k+\frac{1}{2}}$ over \mathbb{C} acts on function $f : \mathbb{H} \rightarrow \mathbb{C}$ by

$$f | \sum_{\nu} c_{\nu} (A_{\nu}, \phi_{\nu}) = \sum_{\nu} c_{\nu} \phi_{\nu}(z)^{-1} f(A_{\nu}z).$$

We require the automorphy factor

$$\phi(z) := \chi(d) \left(\frac{c}{d} \right) \left(\frac{-4}{d} \right)^{-k-\frac{1}{2}} (cz + d)^{k+\frac{1}{2}},$$

where χ is an even Dirichlet character modulo $4p$. As an abuse of notation, we shall let

$$(2.2) \quad A^* := (A, \phi).$$

Let $\Delta_0(4p, \chi)_{k+\frac{1}{2}}$ be the set of these pairs A^* , where $A \in \Gamma_0(4p)$. A meromorphic function $f : \mathbb{H} \rightarrow \mathbb{C}$ is a *weight $k + \frac{1}{2}$ meromorphic modular form with Nebentypus χ* if it is meromorphic at the cusps of $\Gamma_0(4p)$ and if $f|A^* = f$ for all $A^* \in \Delta_0(4p, \chi)_{k+\frac{1}{2}}$.

We let $\mathcal{M}_{k+\frac{1}{2}}(\Gamma_0(4p), \chi)$ be the space of weakly holomorphic modular forms of weight $k + \frac{1}{2}$ with Nebentypus character χ on $\Gamma_0(4p)$. Furthermore, let $\mathcal{M}_{k+\frac{1}{2}}(\Gamma_0(4p))$ denote the case where χ is trivial.

Remark. Although we only describe those forms with level $4p$, most of the following arguments hold for general levels. Furthermore, the facts we quote from works of Shimura and Kohlen were originally stated for holomorphic forms or cusp forms, although their proofs also apply for weakly holomorphic forms.

For $m \geq 1$, define the operators $U(m)$ and $V(m)$ on formal power series in q by

$$(2.3) \quad \left(\sum_{n \in \mathbb{Z}} a(n)q^n \right) |U(m) := \sum_{n \in \mathbb{Z}} a(mn)q^n,$$

$$(2.4) \quad \left(\sum_{n \in \mathbb{Z}} a(n)q^n \right) |V(m) := \sum_{n \in \mathbb{Z}} a(n)q^{nm}.$$

We also require the Atkin-Lehner operators $W(p)$ defined by

$$(2.5) \quad W(p) := \left(\begin{pmatrix} p & a \\ 4p & pb \end{pmatrix}, \left(\frac{-4}{p} \right)^{-k-\frac{1}{2}} p^{-\frac{k}{2}-\frac{1}{4}} (4pz + pb)^{k+\frac{1}{2}} \right),$$

where a and b are integers with $pb - 4a = 1$. The next proposition (see [Sh] and [Ko]) describes the modularity of these operators.

Proposition 2.1. *Suppose that $f \in \mathcal{M}_{k+\frac{1}{2}}(\Gamma_0(4p), \chi)$.*

- (1) *We have $f|U(p), f|W(p) \in \mathcal{M}_{k+\frac{1}{2}}(\Gamma_0(4p), \left(\frac{p}{\cdot}\right)\chi)$.*
- (2) *We have $f|V(p) \in \mathcal{M}_{k+\frac{1}{2}}(\Gamma_0(4p^2), \left(\frac{p}{\cdot}\right)\chi)$.*

We shall employ the trace map $\mathrm{Tr}_4^{4p} : \mathcal{M}_{k+\frac{1}{2}}(\Gamma_0(4p)) \rightarrow \mathcal{M}_{k+\frac{1}{2}}(\Gamma_0(4))$ defined by

$$(2.6) \quad \mathrm{Tr}_4^{4p}(f) := \sum_{i=1}^r f|_{k+\frac{1}{2}} \gamma_i,$$

where $\{\gamma_1, \dots, \gamma_r\}$ is a set of coset representatives for $\Gamma_0(4p) \backslash \Gamma_0(4)$. The next lemma (see page 66 of [Ko]) gives a description of this map in terms of $W(p)$ and $U(p)$.

Lemma 2.2. *If p is an odd prime and $f \in \mathcal{M}_{k+\frac{1}{2}}(\Gamma_0(4p))$, then we have*

$$\mathrm{Tr}_4^{4p}(f) = f + \left(\frac{-1}{p} \right)^{-(k+\frac{1}{2})} \cdot p^{-\frac{k}{2}+\frac{3}{4}} \cdot f|_{k+\frac{1}{2}} W(p)|U(p).$$

3. KOHNEN'S SPACES AND THE TRACES OF SINGULAR MODULI

Let $\mathcal{M}_{k+\frac{1}{2}}^+(\Gamma_0(4p))$ and $\mathcal{M}_{k+\frac{1}{2}}^{+,\epsilon}(\Gamma_0(4p))$ be the spaces defined in (1.3) and (1.4).

3.1. Trace generating functions. Zagier showed how the forms $g_m(z)$ may be used to compute the generating functions for traces and twisted traces of singular moduli. For example, consider

$$(3.1) \quad g_1(z) = q^{-1} + \sum_{D=0}^{\infty} B(1, D)q^D = q^{-1} - 2 + 248q^3 - 492q^4 + \cdots \in \mathcal{M}_{\frac{3}{2}}^+(\Gamma_0(4)).$$

To make the connection to traces, for positive integers m , let

$$(3.2) \quad B_m(1, D) := \text{the coefficient of } q^D \text{ in } g_1(z)|T(m^2),$$

where $T(m^2)$ is the usual Hecke operator on $\mathcal{M}_{\frac{3}{2}}^+(\Gamma_0(4))$. Zagier's formulae for the traces $t_m(D)$ are given by the following theorem (see Theorem 5 of [Za]).

Theorem 3.1. *If $m \geq 1$ and $0 \leq D \equiv 0, 3 \pmod{4}$, then $t_m(D) = -B_m(1, D)$.*

More generally, Zagier (see Theorem 5 of [Za]) obtains the following formula

$$(3.3) \quad B_m(1, D) = \sum_{l|m} lB(l^2, D).$$

Applying Möbius inversion to these formulas immediately gives the following lemma.

Lemma 3.2. *If m is a positive integer and $1 \leq D \equiv 0, 3 \pmod{4}$, then*

$$B(m^2, D) = -\frac{1}{m} \sum_{l|m} \mu(m/l)t_l(D).$$

The following is a special case of Bruinier and Funke's work (see Theorem 1.1 of [BF]).

Theorem 3.3. *If p is an odd prime and $f = \sum a(n)q^n \in \mathcal{M}_0(\Gamma_0^*(p))$, with $a(0) = 0$, then*

$$G_p(f, z) := -\sum_{m \geq 1} \sum_{n \geq 1} ma(-mn)q^{-m^2} + \sum_{n \geq 1} (\sigma_1(n) + p\sigma_1(n/p)) a(-n) + \sum_{D > 0} t_f^*(D)q^D$$

is an element of $\mathcal{M}_{\frac{3}{2}}^{+,+}(\Gamma_0(4p))$, where $\sigma_1(n) := \sum_{t|n} t$ for positive integers n , and $\sigma_1(x) = 0$ for non-integral x .

3.2. Operators on the Kohnen spaces. Here we assume that p is an odd prime. To prove Theorem 1.1, our plan is to apply Lemma 2.2. This requires an explicit description of the action of $U(p)$ and $W(p)$. Although the next two results are proven for holomorphic forms by Kohnen in [Ko], we give their proofs for completeness.

Proposition 3.4. *If $f \in \mathcal{M}_{k+\frac{1}{2}}(\Gamma_0(4p))$, then*

$$f|U(p)|W(p)^2 = \left(\frac{-1}{p}\right)^{k+\frac{1}{2}} f|U(p).$$

Proof. Using (2.1), we obtain the identity

$$W(p)^2 = \left(\frac{-1}{p}\right) \left(\frac{b+1}{4a+pb^2}\right) \left(\frac{-1}{4a+pb^2}\right)^{-k-\frac{1}{2}} \left(\begin{pmatrix} p & 0 \\ 0 & p \end{pmatrix}, 1\right) \\ \left(\begin{pmatrix} p+4a & a(1+b) \\ 4p(1+b) & 4a+pb^2 \end{pmatrix}, \left(\frac{b+1}{4a+pb^2}\right) \left(\frac{-1}{4a+pb^2}\right)^{-k-\frac{1}{2}} ((4p(1+b)z + (4a+pb^2))^{k+\frac{1}{2}})\right).$$

Since $\left(\begin{pmatrix} p & 0 \\ 0 & p \end{pmatrix}, 1\right)$ acts as the identity on $f|U(p)$, and since $\Delta_0(4p, \left(\frac{2}{p}\right)_{k+\frac{1}{2}})$ contains

$$\left(\begin{pmatrix} p+4a & a(1+b) \\ 4p(1+b) & 4a+pb^2 \end{pmatrix}, \left(\frac{b+1}{4a+pb^2}\right) \left(\frac{-1}{4a+pb^2}\right)^{-k-\frac{1}{2}} ((4p(1+b)z + (4a+pb^2))^{k+\frac{1}{2}})\right),$$

Proposition 2.1 gives the identity

$$f|U(p)|W(p)^2 = \left(\frac{-1}{p}\right) \left(\frac{b+1}{4a+pb^2}\right) \left(\frac{-1}{4a+pb^2}\right)^{-k-\frac{1}{2}} f|U(p).$$

Since $\left(\frac{-1}{4a+pb^2}\right) = \left(\frac{-1}{p}\right)$, to complete the proof we have to show that $\left(\frac{b+1}{4a+pb^2}\right) = 1$. For this we choose $l \in \mathbb{N}$ maximal such that $2^l | (b+1)$. Moreover, we assume that $a \equiv 0 \pmod{2}$ (i.e, $b \equiv p \pmod{8}$). The Law of Quadratic Reciprocity implies that

$$(3.4) \quad \left(\frac{b+1}{4a+pb^2}\right) = \left(\frac{2}{p}\right)^l (-1)^{\frac{1}{4}(\frac{b+1}{2^l}-1)(p-1)} \left(\frac{-1}{\left(\frac{b+1}{2^l}\right)}\right).$$

Here we used the fact that $4a - pb = -1$. To prove that the right-hand side of (3.4) equals 1, we distinguish the cases whether $l > 1$ or $l = 1$. If $l > 1$, then $p \equiv b \equiv 3 \pmod{4}$. Using that in this case $(-1)^{\frac{1}{4}(\frac{b+1}{2^l}-1)(p-1)} = \left(\frac{-1}{(b+1)/2^l}\right)$, we obtain from (3.4) that $\left(\frac{b+1}{4a+pb^2}\right) = \left(\frac{2}{p}\right)^l$. Thus the claim is clear if l is even. If l is odd, then $p \equiv b \equiv -1 \pmod{8}$ and therefore $\left(\frac{2}{p}\right) = 1$. Thus we also obtain the claim. If $l = 1$, then $p \equiv b \equiv 1, -3 \pmod{8}$. If $p \equiv b \equiv 1 \pmod{8}$, then we obtain from (3.4) that $\left(\frac{b+1}{4a+pb^2}\right) = 1$, since $\left(\frac{2}{p}\right) = 1$, $(-1)^{\frac{1}{4}(\frac{b+1}{2}-1)(p-1)} = 1$, and $\left(\frac{-1}{(b+1)/2}\right) = 1$. If $p \equiv b \equiv 5 \pmod{8}$, we also obtain $\left(\frac{b+1}{4a+pb^2}\right) = 1$ from (3.4), since $\left(\frac{2}{p}\right) = -1$, $(-1)^{\frac{1}{4}(\frac{b+1}{2}-1)(p-1)} = 1$, and $\left(\frac{-1}{(b+1)/2}\right) = -1$. \square

The following lemma relates the action of $U(p)$ and $W(p)$ on $\mathcal{M}_{k+\frac{1}{2}}^{+, \epsilon}(\Gamma_0(4p))$.

Lemma 3.5. *If $f \in \mathcal{M}_{k+\frac{1}{2}}^{+, \epsilon}(\Gamma_0(4p))$, where $\epsilon \in \{\pm 1\}$, then $f|p^{-\frac{k}{2}+\frac{1}{4}}U(p)|W(p)$ is in $\mathcal{M}_{k+\frac{1}{2}}^{+, \epsilon}(\Gamma_0(4p))$. Moreover, we have*

$$f|W(p) = \epsilon \left(\frac{-1}{p}\right)^{k+\frac{1}{2}} p^{-\frac{k}{2}+\frac{1}{4}} f|U(p).$$

Proof. Exactly as in [Ko] p. 41 we obtain for $f = \sum_{n \gg -\infty} a(n)q^n$ the following identity

$$(3.5) \quad f|p^{-\frac{k}{2}+\frac{1}{4}}U(p)|W(p) = \sum_{n \gg -\infty} \left(\frac{(-1)^{kn}}{p} \right) a(n)q^n + p^{-\frac{1}{2}}f| \left(\begin{pmatrix} 1 & \nu_0 \\ 0 & p \end{pmatrix}, p^{\frac{k}{2}+\frac{1}{4}} \right) |W(p),$$

where ν_0 is an integer with $1 + 4\nu_0 \equiv 0 \pmod{p}$. Since $f \in \mathcal{M}_{k+\frac{1}{2}}^{+, \epsilon}(\Gamma_0(4p))$, it is clear that the Fourier coefficients of the first term vanish unless $\left(\frac{(-1)^{kn}}{p}\right) = \epsilon$. Moreover, the coefficients of the second term vanish unless $p|n$ which follows directly from the identity

$$(3.6) \quad \left(\begin{pmatrix} 1 & \nu_0 \\ 0 & p \end{pmatrix}, p^{\frac{k}{2}+\frac{1}{4}} \right) W(p) = \left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \left(\frac{-1}{p} \right)^{k+\frac{1}{2}} \right) C^* W(p) \left(\begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix}, p^{-\frac{k}{2}-\frac{1}{4}} \right),$$

where $C \in \Gamma_0(4p)$. Since Kohnen (see page 39 of [Ko]) proved that

$$f|p^{-\frac{k}{2}+\frac{1}{4}}U(p)|W(p) \in \mathcal{M}_{k+\frac{1}{2}}(\Gamma_0(4p)),$$

these calculations show that $f|p^{-\frac{k}{2}+\frac{1}{4}}U(p)|W(p) \in \mathcal{M}_{k+\frac{1}{2}}^{+, \epsilon}(\Gamma_0(4p))$.

For the second part of the lemma, note that (3.5) and (3.6) imply that

$$f|p^{-\frac{k}{2}+\frac{1}{4}}U(p)|W(p) = \epsilon f - \epsilon f|U(p)|V(p) + \left(\frac{-1}{p} \right)^{-k-\frac{1}{2}} p^{\frac{k}{2}-\frac{1}{4}} f|W(p)|V(p).$$

Since $f|p^{-\frac{k}{2}+\frac{1}{4}}U(p)|W(p) \in \mathcal{M}_{k+\frac{1}{2}}^{+, \epsilon}(\Gamma_0(4p))$, we have

$$h^* := -\epsilon f|U(p)|V(p) + \left(\frac{-1}{p} \right)^{-k-\frac{1}{2}} p^{\frac{k}{2}-\frac{1}{4}} f|W(p)|V(p) \in \mathcal{M}_{k+\frac{1}{2}}^{+, \epsilon}(\Gamma_0(4p)).$$

If $h^* = \sum_{n \gg -\infty} c(n)q^n$, then the coefficients $c(n)$ vanish unless $p|n$. A theorem of Serre and Stark (Theorem 1 of [SS]) then implies that $h^* = h_0|V(p)$, where $h_0 \in \mathcal{M}_{k+\frac{1}{2}}(\Gamma_0(4), \left(\frac{\epsilon}{p}\right))$. Obviously, there are no such non-zero forms, and so $h_0 = 0$ which in turn implies that

$$f|p^{-\frac{k}{2}+\frac{1}{4}}U(p)|W(p) = \epsilon f.$$

The second claim in the lemma now follows by applying Proposition 3.4. \square

Remark. The theorem of Serre and Stark employed above is stated for holomorphic half-integral weight forms. Note that we may apply their result to $h^*(z)\Delta(pz)^a$, where $\Delta(z)$ is the usual normalized weight 12 cusp form on $SL_2(\mathbb{Z})$ and a is a sufficiently large positive integer. This results in the desired conclusion that $h^* = 0$.

3.3. Proof of Theorem 1.1. Now we prove Theorem 1.1 using the results from the previous two sub-sections.

Proof of Theorem 1.1. Lemma 2.2 and Lemma 3.5 imply that

$$\mathrm{Tr}_4^{4p}(g) = g + \left(\frac{-1}{p} \right)^{-\frac{3}{2}} p^{\frac{1}{4}} \cdot g|W(p)|U(p) = g + \epsilon g|U(p^2).$$

Therefore, we find that

$$\mathrm{Tr}_4^{4p}(g) = \sum_{m \geq 1} (a(-m) + \epsilon a(-mp^2)) q^{-m} + \sum_{n \geq 0} (a(n) + \epsilon a(np^2)) q^n.$$

By its Fourier expansion and Lemma 2.2, we have that $\mathrm{Tr}_4^{4p}(g) \in \mathcal{M}_{\frac{3}{2}}^+(\Gamma_0(4))$, and is given by

$$\mathrm{Tr}_4^{4p}(g) = \sum_{m \geq 1} (a(-m) + \epsilon a(-mp^2)) g_m(z).$$

Here we require that the g_m are uniquely determined by their “principal parts”, together with the fact that there are no holomorphic forms in $\mathcal{M}_{\frac{3}{2}}^+(\Gamma_0(4))$. \square

Proof of Corollary 1.2. By applying Theorem 1.1 to the modular form $G_p(f, z)$ from Theorem 3.3, we obtain the identity

$$\mathrm{Tr}_4^{4p}(G_p(f, z)) = - \sum_{m \geq 1} \left(\sum_{n \geq 1} ma(-mn) + p^2 ma(-mp^2 n) \right) g_{m^2}(z).$$

The corollary now follows from Theorem 3.3 and Lemma 3.2. \square

REFERENCES

- [BF] J.H. Bruinier and J. Funke, Traces of CM values of modular forms, preprint.
- [Ko] W. Kohnen, Newforms of half-integral weight, *J. Reine Angew. Math.* **333**, (1982) pages 32–72.
- [Sh] G. Shimura, On modular forms of half integral weight, *Ann. of Math. (2)* **97**, (1973) pages 440–481.
- [SS] J.-P. Serre and H. M. Stark, Modular forms of weight 1/2. Modular functions of one variable, VI (Proc. Second Internat. Conf., Univ. Bonn, Bonn, 1976), pp. 27–67. Lecture Notes in Math., Vol. 627, Springer, Berlin, 1977.
- [Za] D. Zagier, Traces of singular moduli, Motives, polylogarithms and Hodge theory, Part I (Irvine, CA, 1998), (2002), Int. Press Lect. Ser., 3, I, Int. Press, Somerville, MA, pages 211–244.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN, MADISON, WISCONSIN 53706
E-mail address: bringman@math.wisc.edu
E-mail address: ono@math.wisc.edu