

# DIVISIBILITY CRITERIA FOR CLASS NUMBERS OF IMAGINARY QUADRATIC FIELDS

PAUL JENKINS AND KEN ONO

ABSTRACT. In a recent paper, Guerzhoy obtained formulas for certain class numbers as  $p$ -adic limits of traces of singular moduli. Using earlier work by Bruinier and the second author, we derive a more precise form of these results using results of Zagier. Specifically, if  $-d < -4$  is a fundamental discriminant and  $n$  is a positive integer, then

$$\mathrm{Tr}(p^{2n}d) \equiv \frac{24}{p-1} \cdot \left(1 - \left(\frac{-d}{p}\right)\right) \cdot H(-d) \pmod{p^n}$$

provided that  $p \in \{2, 3\}$  and  $\left(\frac{-d}{p}\right) = -1$ , or  $p \in \{5, 7, 13\}$  and  $\left(\frac{-d}{p}\right) \neq 1$ .

## 1. INTRODUCTION AND STATEMENT OF RESULTS

Throughout, let  $d \equiv 0, 3 \pmod{4}$  be a positive integer, and let  $\mathcal{Q}_d$  denote the set of positive definite integral binary quadratic forms  $Q(x, y) = ax^2 + bxy + cy^2 = [a, b, c]$  with discriminant  $-d = b^2 - 4ac$  (including imprimitive forms if there are any). The group  $\Gamma := \mathrm{PSL}_2(\mathbb{Z})$  acts on  $\mathcal{Q}_d$  with finitely many orbits, and if  $\omega_Q$  is defined by

$$\omega_Q = \begin{cases} 2 & \text{if } Q \sim_{\Gamma} [a, 0, a], \\ 3 & \text{if } Q \sim_{\Gamma} [a, a, a], \\ 1 & \text{otherwise,} \end{cases}$$

then the Hurwitz-Kronecker class number  $H(-d)$  is given by

$$(1.1) \quad H(d) = \sum_{Q \in \mathcal{Q}_d/\Gamma} \frac{1}{\omega_Q}.$$

If  $-d < -4$  is a fundamental discriminant, then  $H(-d)$  is the class number of the ring of integers of the imaginary quadratic field  $\mathbb{Q}(\sqrt{-d})$ .

Recently, Guerzhoy has obtained some interesting expressions for  $\left(1 - \left(\frac{-d}{p}\right)\right) H(-d)$  as  $p$ -adic limits of traces of singular moduli. To make this precise, we first recall some notation. For positive definite binary quadratic forms  $Q$ , let  $\alpha_Q$  be the unique root of

---

*Date:* August 31, 2006.

*2000 Mathematics Subject Classification.* 11R29, 11F37.

*Key words and phrases.* Class numbers, singular moduli.

The authors thank the National Science Foundation for their generous support. The second author is grateful for the support of a Packard and a Romnes Fellowship.

$Q(x, 1) = 0$  in the upper half of the complex plane. If  $j(z)$  is the usual  $\mathrm{SL}_2(\mathbb{Z})$  modular function

$$j(z) = \frac{E_4(z)^3}{\Delta(z)} = q^{-1} + 744 + 196884q + \cdots,$$

where  $q = e^{2\pi iz}$ , then define integers  $\mathrm{Tr}(d)$  by

$$(1.2) \quad \mathrm{Tr}(d) = \sum_{Q \in \mathcal{Q}_d/\Gamma} \frac{j(\alpha_Q) - 744}{\omega_Q}.$$

The algebraic integers  $j(\alpha_Q)$  are known as *singular moduli*. Guerzhoy proved (see Corollary 2.4 (a) of [5]) that if  $p \in \{3, 5, 7, 13\}$  and  $-d < -4$  is a fundamental discriminant, then one has the  $p$ -adic limit formula

$$(1.3) \quad \left(1 - \left(\frac{-d}{p}\right)\right) \cdot H(-d) = \frac{p-1}{24} \lim_{n \rightarrow +\infty} \mathrm{Tr}(p^{2n}d).$$

If  $\left(\frac{-d}{p}\right) = 1$ , then this result simply implies that  $\mathrm{Tr}(p^{2n}d) \rightarrow 0$   $p$ -adically as  $n$  tends to infinity. Thanks to work of Boylan, Edixhoven and the first author (see [2, 4, 6]), it turns out that more is true. In particular, if  $p$  is any prime and  $\left(\frac{-d}{p}\right) = 1$ , then

$$(1.4) \quad \mathrm{Tr}(p^{2n}d) \equiv 0 \pmod{p^n}.$$

In earlier work [3], Bruinier and the second author obtained certain  $p$ -adic expansions for  $H(-d)$  in terms of the Borcherds exponents of certain modular functions with Heegner divisor. In his paper [5], Guerzhoy asks whether there is a connection between (1.3) and these results when  $\left(\frac{-d}{p}\right) \neq 1$ . In this note we show that this is indeed the case by establishing the following congruences.

**Theorem 1.1.** *Suppose that  $-d < -4$  is a fundamental discriminant and that  $n$  is a positive integer. If  $p \in \{2, 3\}$  and  $\left(\frac{-d}{p}\right) = -1$ , or  $p \in \{5, 7, 13\}$  and  $\left(\frac{-d}{p}\right) \neq 1$ , then*

$$\frac{24}{p-1} \cdot \left(1 - \left(\frac{-d}{p}\right)\right) \cdot H(-d) \equiv \mathrm{Tr}(p^{2n}d) \pmod{p^n}.$$

*In particular, under these hypotheses  $p^n$  divides  $\frac{24}{p-1} \left(1 - \left(\frac{-d}{p}\right)\right) \cdot H(-d)$  if and only if  $p^n$  divides  $\mathrm{Tr}(p^{2n}d)$ .*

*Three remarks.*

1) Theorem 1.1 includes  $p = 2$ . For simplicity, Guerzhoy chose to work with odd primes  $p$ , and this explains the omission of  $p = 2$  in (1.3).

2) Despite the uniformity of (1.4), it turns out that the restriction on  $p$  in Theorem 1.1 is required. For example, if  $p = 11$ ,  $n = 1$  and  $-d = -15$ , then  $\left(\frac{-15}{11}\right) = -1$ ,  $H(-15) = 2$ ,

and we have

$$\begin{aligned} \text{Tr}(11^2 \cdot 15) &= -13374447806956269126908865521582974841084501554961922745794 \\ &\equiv 7 \not\equiv \frac{48}{10} \cdot H(-15) \pmod{11}. \end{aligned}$$

3) There are generalizations of Theorem 1.1 which hold for primes  $p \notin \{2, 3, 5, 7, 13\}$ . For example, one may employ Serre's theory [7] of  $p$ -adic modular forms to derive more precise versions of Corollary 2.4 (b) of [5].

## 2. THE PROOF OF THEOREM 1.1

The proof of Theorem 1.1 follows by combining earlier work of Bruinier and the second author with results of Zagier and a combinatorial formula used earlier by the first author. We recall some necessary notation.

Let  $M_{\lambda+\frac{1}{2}}^!$  be the space of weight  $\lambda + \frac{1}{2}$  weakly holomorphic modular forms on  $\Gamma_0(4)$  with Fourier expansion

$$f(z) = \sum_{(-1)^\lambda n \equiv 0,1 \pmod{4}} a(n)q^n.$$

For  $0 \leq d \equiv 0, 3 \pmod{4}$ , we let  $f_d(z)$  be the unique form in  $M_{\frac{1}{2}}^!$  with expansion

$$(2.1) \quad f_d(z) = q^{-d} + \sum_{0 < D \equiv 0,1 \pmod{4}} A(D, d)q^D.$$

The coefficients  $A(D, d)$  of the  $f_d$  are integers. For completeness, we set  $A(M, N) = 0$  if  $M$  or  $N$  is not an integer. These modular forms are described in detail in [8].

For fundamental discriminants  $-d < -4$ , Borcherds' theory on the infinite product expansion of modular forms with Heegner divisor [1] implies that

$$q^{-H(-d)} \prod_{n=1}^{\infty} (1 - q^n)^{A(n^2, d)}$$

is a weight zero modular function on  $\text{SL}_2(\mathbb{Z})$  whose divisor consists of a pole of order  $H(-d)$  at infinity and a simple zero at each Heegner point of discriminant  $-d$ . Using this factorization, Bruinier and the second author proved the following theorem.

**Theorem 2.1** ([3], Corollary 3). *Let  $-d < -4$  be a fundamental discriminant. If  $p \in \{2, 3\}$  and  $\left(\frac{-d}{p}\right) = -1$ , or  $p \in \{5, 7, 13\}$  and  $\left(\frac{-d}{p}\right) \neq 1$ , then as  $p$ -adic numbers we have*

$$H(-d) = \frac{p-1}{24} \sum_{k=0}^{\infty} p^k A(p^{2k}, d).$$

*Remark.* The case when  $p = 13$  is not proven in [3]. However, thanks to the remark preceding Theorem 8 of [7] on 13-adic modular forms with weight congruent to 2 (mod 12), and Theorem 2 of [3], the proof of Corollary 3 [3] still applies *mutatis mutandis*.

Zagier identified traces of singular moduli with the coefficients  $A(D, d)$  as follows.

**Theorem 2.2** ([8], Corollary to Theorem 3). *For all positive integers  $d \equiv 0, 3 \pmod{4}$ ,*

$$\mathrm{Tr}(d) = A(1, d).$$

Combining Zagier's duality ([8], Theorem 4) between coefficients of modular forms in  $M_{\frac{1}{2}}^1$  and in  $M_{\frac{3}{2}}^1$  with the action of the Hecke operators on these spaces, the first author proved the following combinatorial formula.

**Lemma 2.3** ([6], Theorem 1.1). *If  $p$  is a prime and  $d, D, n$  are positive integers such that  $-d, D \equiv 0, 1 \pmod{4}$ , then*

$$\begin{aligned} A(D, p^{2n}d) = & p^n A(p^{2n}D, d) + \sum_{k=0}^{n-1} \left(\frac{D}{p}\right)^{n-k-1} \left( A\left(\frac{D}{p^2}, p^{2k}d\right) - p^{k+1} A\left(p^{2k}D, \frac{d}{p^2}\right) \right) \\ & + \sum_{k=0}^{n-1} \left(\frac{D}{p}\right)^{n-k-1} \left( \left( \left(\frac{D}{p}\right) - \left(\frac{-d}{p}\right) \right) p^k A(p^{2k}D, d) \right). \end{aligned}$$

*Remark.* This result is stated for odd  $p$  in [6], but the proof holds for  $p = 2$  as well.

*Proof of Theorem 1.1.* Under the given hypotheses, Theorem 2.1 implies that

$$(2.2) \quad \frac{24}{p-1} \cdot H(-d) \equiv \sum_{k=0}^{n-1} p^k A(p^{2k}, d) \pmod{p^n}.$$

By letting  $D = 1$  in Lemma 2.3, for these  $d$  and  $p$  we find that

$$(2.3) \quad \left(1 - \left(\frac{-d}{p}\right)\right) \sum_{k=0}^{n-1} p^k A(p^{2k}, d) = A(1, p^{2n}d) - p^n A(p^{2n}, d).$$

Inserting this expression for the sum into (2.2), we conclude that

$$\frac{24}{p-1} \cdot \left(1 - \left(\frac{-d}{p}\right)\right) \cdot H(-d) \equiv A(1, p^{2n}d) \pmod{p^n},$$

which by Zagier's theorem is  $\mathrm{Tr}(p^{2n}d)$ . □

## REFERENCES

- [1] R. E. Borcherds, *Automorphic forms on  $O_{s+2,2}(\mathbb{R})$  and infinite products*, Invent. Math. **120** (1995), no. 1, 161–213.
- [2] M. Boylan, *2-adic properties of Hecke traces of singular moduli*, Math. Res. Lett. **12** (2005), no. 4, 593–609.

- [3] J. H. Bruinier and K. Ono, *The arithmetic of Borchers' exponents*, Math. Ann. **327** (2003), no. 2, 293–303.
- [4] B. Edixhoven, *On the  $p$ -adic geometry of traces of singular moduli*, International Journal of Number Theory **1** (2005), 495–497.
- [5] P. Guerzhoy, *The Borchers-Zagier isomorphism and a  $p$ -adic version of the Kohnen-Shimura map*, Int. Math. Res. Not. (2005), no. 13, 799–814.
- [6] P. Jenkins,  *$p$ -adic properties for traces of singular moduli*, International Journal of Number Theory **1** (2005), 103–107.
- [7] J.-P. Serre, *Formes modulaires et fonctions zêta  $p$ -adiques*, Modular functions of one variable, III (Proc. Internat. Summer School, Univ. Antwerp, 1972), Springer, Berlin, 1973, pp. 191–268. Lecture Notes in Math., Vol. 350.
- [8] D. Zagier, *Traces of singular moduli*, Motives, polylogarithms and Hodge theory, Part I (Irvine, CA, 1998), Int. Press Lect. Ser., vol. 3, Int. Press, Somerville, MA, 2002, pp. 211–244.

UCLA MATHEMATICS DEPARTMENT, LOS ANGELES, CALIFORNIA 90095-1555  
*E-mail address:* `jenkins@math.ucla.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN, MADISON, WISCONSIN 53706  
*E-mail address:* `ono@math.wisc.edu`