

A FEW BRST BICOMPLEXES

Rudolf Schmid*
Department of Mathematics
Emory University
Atlanta, Georgia 30322

Abstract

There are many double complexes in the mathematics and physics literature which are related to BRST transformations and anomalies, e.g. Variational-, BRST-, Faddeev-, Koszul-Tate-, Weil-, Gelfand-Fuks-, semi infinite-, Cech-DeRham-, foliation-, Lie group/algebra-bicomplex. The goal is to identify them and to establish relations between them; to compute their cohomologies and the corresponding anomalies. Here we list and identify only few of them and refer to [11] for more details.

Lecture given at 21st International Conference on Differential Geometry Methods in Theoretical Physics, June 5-9, 1992, Nankai Institute of Mathematics, Tianjing, P.R. China.

*Supported in part by Emory University Research Grant # 2-31220

1 Variational bicomplex

Of all the bicomplexes mentioned above, the variational bicomplex is the "smallest" in the following sense. Let $\pi : P \rightarrow M^n$ be a fiber bundle and consider the jet bundle $J^\infty(\pi)$. Its exterior forms are bigraded by horizontal (r) and vertical (s) degrees

$$\Omega^p(J^\infty(\pi)) = \bigoplus_{r+s=p} \Omega^{r,s}(J^\infty(\pi)), \quad d = d_H + d_V : \Omega^{r,s} \rightarrow \Omega^{r+1,s} \oplus \Omega^{r,s+1}.$$

Horizontal and vertical derivatives satisfy Poincare lemmas and $d_H^2 = 0$, $d_V^2 = 0$, $d_H d_V = -d_V d_H$, hence $d^2 = 0$. With $\mathcal{F}^s = \{\omega \in \Omega^{n,s} | I(\omega) = \omega\}$ (functional forms), I the Euler operator we have the augmented variational bicomplex [1]:

$$\begin{array}{ccccccccccc}
 & & \uparrow d_V & & & & \uparrow d_V & & \uparrow d_V & & \\
 0 & \longrightarrow & \Omega^{0,3} & & \dots & & \Omega^{n,3} & \xrightarrow{I} & \mathcal{F}^3 & \longrightarrow & 0 \\
 & & \uparrow d_V & & & & \uparrow d_V & & \uparrow d_V & & \\
 0 & \longrightarrow & \Omega^{0,2} & \xrightarrow{d_H} & \Omega^{1,2} & \xrightarrow{d_H} \dots & \Omega^{n-1,2} & \xrightarrow{d_H} & \Omega^{n,2} & \xrightarrow{I} & \mathcal{F}^2 & \longrightarrow & 0 \\
 & & \uparrow d_V & & \uparrow d_V & & \uparrow d_V & & \uparrow d_V & & \uparrow d_V & & \\
 0 & \longrightarrow & \Omega^{0,1} & \xrightarrow{d_H} & \Omega^{1,1} & \xrightarrow{d_H} \dots & \Omega^{n-1,1} & \xrightarrow{d_H} & \Omega^{n,1} & \xrightarrow{I} & \mathcal{F}^1 & \longrightarrow & 0 \\
 & & \uparrow d_V & & \uparrow d_V & & \uparrow d_V & & \uparrow d_V & & & & \\
 \mathbf{R} & \longrightarrow & \Omega^{0,0} & \xrightarrow{d_H} & \Omega^{1,0} & \xrightarrow{d_H} \dots & \Omega^{n-1,0} & \xrightarrow{d_H} & \Omega^{n,0} & & & &
 \end{array}$$

The *local* cohomologies $H^p(\pi) = \ker d / \text{im } d$ can be computed explicitly using spectral sequences [1]. These are *local* cohomologies because for any $\omega \in \Omega^{r,s}(J^\infty(\pi))$ we have $\omega(j, X_1, \dots, X_s) \in \Omega^r(M)$, for any jet $j \in J^\infty(\pi)$ and vector fields X_i on $J^\infty(\pi)$.

2 BRST bicomplex

The BRST bicomplex described in [2],[9],[10] is related to the variational bicomplex as follows: Let $\pi : P \rightarrow M$ be a principal G -bundle and $\pi^p : \Omega^p(P, \text{Lie } G) \rightarrow M$ Lie algebra valued equivariant p-forms. Let \mathcal{G} denote the Lie group of gauge transformations and $\text{Lie } \mathcal{G}$ its Lie algebra. Set $\mathbf{C}_{loc}^{q,p} =$

$\Lambda^q(Lie\mathcal{G}, \Omega^{p,0}(J^\infty(\pi^P)))$ (local q - cochains) and define $\delta_{loc} : \mathbf{C}_{loc}^{q,p} \rightarrow \mathbf{C}_{loc}^{q+1,p}$ to be the Chevalley-Eilenberg coboundary operator with respect to a representation ρ :

$$\begin{aligned} (\delta_{loc}\phi)(\xi_0, \dots, \xi_q) &= \sum_{i=0}^q (-1)^i \rho'(\xi_i) \phi(\xi_0, \dots, \hat{\xi}_i, \dots, \xi_q) \\ &+ \sum_{i<j} (-1)^{i+j} \phi(\rho'(\xi_i)\xi_j, \dots, \hat{\xi}_i, \dots, \hat{\xi}_j, \dots, \xi_q), \end{aligned}$$

where ρ' is the derived representation of $Lie\mathcal{G}$ on $\Omega(P, LieG)$ induced by a representation ρ of G . We have $\delta_{loc}^2 = 0$. Then we define the BRST operator $\mathbf{s} : \mathbf{C}_{loc}^{q,p} \rightarrow \mathbf{C}_{loc}^{q+1,p}$ as

$$\mathbf{s} \equiv \frac{(-1)^{p+1}}{q+1} \delta_{loc} .$$

It is clear that \mathbf{s} is nilpotent, $\mathbf{s}^2 = 0$. We call the associated *local* cohomology of this bicomplex $\{\mathbf{C}_{loc}^{q,p}, \mathbf{s}\}$ the *BRST cohomology* of the gauge algebra $Lie\mathcal{G}$, denoted by $H_{BRST}^*(Lie\mathcal{G})$. In [9],[10] we derived the classical BRST transformations using the Chevalley-Eilenberg construction for the *adjoint* representation:

$$\mathbf{s}A = d\eta + [A, \eta] , \quad \mathbf{s}\eta = -\frac{1}{2}[\eta, \eta] , \quad \mathbf{s}\bar{\eta} = b , \quad \mathbf{s}b = 0 ,$$

where the vector potential $A \in \mathbf{C}_{loc}^{0,1}$ and the ghost field $\eta \in \mathbf{C}_{loc}^{1,0}$ is the Maurer Cartan form on \mathcal{G} . We derive a homotopy formula on this bicomplex and with the introduction of Chern-Simons type forms $\omega_{2q-i}^{i-1} = a_i p(A, [A, A]^{i-1}, F_A^{q-1})$ we obtain the associated descent equations $\delta\omega_{2q-1}^0 = -d\omega_{2q-2}^1$, $\delta\omega_{2q-2}^1 = -d\omega_{2q-3}^2$, . . . , $\delta\omega_0^{2q-1} = 0$. We identify the non-Abelian anomaly, which automatically satisfies the Wess-Zumino consistency condition, as a cohomology class in $H_{loc}^1(Lie\mathcal{G})$ represented by ω_{2q-2}^1 in $n = 2q-2$ dimensions.

For example, for $q = 2, q = 3$ we get the 2- and 4-dimensional non-Abelian anomaly respectively, represented by $\omega_2^1 = Tr(\eta\delta_{loc}\tilde{A})$, and $\omega_4^1 = Tr(\eta\delta_{loc}(\tilde{A}\delta_{loc}\tilde{A} + \frac{2}{3}\tilde{A}^3))$ resp., where $\tilde{A} = A + \eta$.

3 Faddeev's bicomplex

Let $\pi : (P, G) \rightarrow M$ be a principal bundle and consider $G^p = \{f : S^p \rightarrow G | \infty \rightarrow 1\}$, the space of p -loops. We have the exterior derivative $d : \Omega^q(P \times G^p) \rightarrow \Omega^{q+1}(P \times G^p)$ and the simplicial group coboundary operator $\Delta : \Omega^{q-p}(P \times G^p) \rightarrow \Omega^{q-p}(P \times G^{p+1})$ induced by $\Delta_i : P \times G^{p+1} \rightarrow P \times G^p : (x, g_1, \dots, g_{p+1}) \mapsto (x, g_1, \dots, g_i g_{i+1}, \dots, g_{p+1})$.

For example for S^3 the Chern-Simon form is $\omega_5^0 = Tp(A)$, where $dTp(A) = p(F) = Trace F^3$ (p =invar. polynomial, T = transgression). We get the stair case equations [12]:

$$\begin{array}{c|cccc}
 q=3 & \text{Tr } F^3 & & & \\
 & \uparrow d & & & \\
 q=2 & \omega_5^0 & \xrightarrow{\Delta} & \bullet & \\
 & & & \uparrow d & \\
 q=1 & & & \omega_4^1 & \xrightarrow{\Delta} & \bullet \\
 & & & & & \uparrow d \\
 q=0 & & & & & \omega_3^2 & \xrightarrow{\Delta} & 0 \\
 & & & & & p=2 & & p=3 \\
 & & & & & p=0 & & p=1
 \end{array}$$

ω_3^2 represents the anomaly.

4 Koszul-Tate complex

Let M be a Poisson manifold with a Hamiltonian G action. Extend the momentum map $J : LieG \rightarrow C^\infty(M)$ to a super derivative δ and extend the Lie algebra d , $d : \Lambda LieG \otimes C^\infty(M) \rightarrow Lie^*G \otimes (\Lambda LieG \otimes C^\infty(M))$ defined by $dk(\xi) = \xi \cdot k$, (\cdot = repres. of $LieG$ on $\Lambda LieG \otimes C^\infty(M)$) to \tilde{d} such that we have

$$\begin{array}{ccc}
\Lambda^p \text{Lie}^* G \otimes \Lambda^q \text{Lie} G \otimes C^\infty(M) & \xrightarrow{\delta} & \Lambda^p \text{Lie}^* G \otimes \Lambda^{q-1} \text{Lie} G \otimes C^\infty(M) \\
& \tilde{d} \downarrow & \\
\Lambda^{p+1} \text{Lie}^* G \otimes \Lambda^q \text{Lie} G \otimes C^\infty(M) & &
\end{array}$$

δ and \tilde{d} being defined by

$$\delta(\omega \otimes \xi \otimes 1) = \omega \otimes 1 \otimes J(\xi), \quad \delta(\omega \otimes 1 \otimes f) = \omega \otimes 0, \quad \tilde{d}(\omega \otimes k) = d\omega \otimes k + (-1)^p \wedge dk.$$

We have $\delta^2 = 0$, $\tilde{d}^2 = 0$ and $\tilde{d}\delta = \delta\tilde{d}$. The total differential defines the BRST operator $D = \tilde{d} + (-1)^p 2\delta : \mathcal{C}^k \rightarrow \mathcal{C}^{k+1}$, satisfying nilpotency $D^2 = 0$, where $\mathcal{C}^k = \sum_{p-q=k} \Lambda^p \text{Lie}^* G \otimes \Lambda^q \text{Lie} G \otimes C^\infty(M)$.

The functions on the reduced phase space are given by the cohomology [8]

$$C^\infty(J^{-1}(0)/G) = H_D(\Lambda \text{Lie}^* G \otimes \Lambda \text{Lie} G \otimes C^\infty(M)).$$

which equals the space $E_2^{0,0}$ of the associated spectral sequence.

5 Weil complex

Let $\Lambda(\text{Lie}^* \mathcal{G})$ be the exterior algebra and $\mathbf{S}(\text{Lie}^* \mathcal{G})$ the symmetric algebra of $\text{Lie} \mathcal{G}$, the Lie algebra of infinitesimal gauge transformations. The Weil algebra $W(\text{Lie} \mathcal{G}) \equiv \Lambda(\text{Lie}^* \mathcal{G}) \otimes \mathbf{S}(\text{Lie}^* \mathcal{G})$ is a graded differential \mathcal{G} algebra

$$W(\text{Lie} \mathcal{G}) = \sum_r W^r, \quad W^r = \sum_{p+2q=r} \Lambda^p(\text{Lie}^* \mathcal{G}) \otimes S^q(\text{Lie}^* \mathcal{G}).$$

Let $\{e_\alpha\}$ be a basis of $\text{Lie} \mathcal{G}$ and $\{\theta^\alpha\}$ its dual basis of $\text{Lie}^* \mathcal{G}$, and let $\{u^\alpha\}$ be a basis of $\mathbf{S}(\text{Lie}^* \mathcal{G})$. The antiderivation δ_W of degree 1 on $W(\text{Lie} \mathcal{G})$ is given by $\delta_W = \delta_\Lambda + \delta_S + h$, where $\delta_\Lambda : W^p \rightarrow W^{p+1}$ is given by: for $\phi \in \Lambda^p(\text{Lie} \mathcal{G})$, $x_i \in \text{Lie} \mathcal{G}$

$$(\delta_\Lambda \phi)(x_0, \dots, x_p) = \sum_{\nu < \mu} (-1)^{\nu+\mu} \phi([x_\nu, x_\mu], x_0, \dots, \hat{x}_\nu, \dots, \hat{x}_\mu, \dots, x_p),$$

or $\delta_\Lambda = \frac{1}{2} \sum_\alpha \mu(\theta^\alpha) L_\Lambda(e_\alpha)$, where L_Λ is the Lie derivative and $\mu(a)b \equiv a \wedge b$. We have $\delta_\Lambda^2 = 0$. Moreover $\delta_S = \sum_\alpha \mu(\theta^\alpha) L_S(e_\alpha) : W^p \rightarrow W^{p+1}$. Note that

$\delta_s^2 \neq 0$ but $(\delta_\Lambda + \delta_S)^2 = 0$, so $\delta_S^2 = -(\delta_\Lambda \delta_S + \delta_S \delta_\Lambda)$. The operator $h : W^p \rightarrow W^{p+1}$ is defined by $h = \sum_\alpha \mu_S(\theta^\alpha) \otimes i_A(e_\alpha)$ and is an antiderivative of degree 1 ($i_A =$ interior product). The BRST operator is the total differential $\delta_W = \delta_\Lambda + \delta_S + h$. The associated anomalies in $H^1(\text{Lie}\mathcal{G})$ can be computed explicitly [7].

References

- [1] I.M. Anderson, *The Variational Bicomplex*, preprint, 1991.
- [2] L. Bonora, P. Cotta-Ramusino, *Some Remarks on BRS Transformations, Anomalies and the Cohomology of the Lie Group of Gauge Transformations*, Comm. Math. Phys. **87** (1983), 589-603.
- [3] J.A. Dixon, *Calculations of BRS Cohomology with Spectral Sequences*, Comm. Math. Phys. **139** (1991), 495-526.
- [4] M. Dubois-Violette, *The Weil-B.R.S. Algebra of a Lie Algebra and the Anomalous Term in Gauge Theories*, J. Geom. Phys. **3** (1986) 525-565.
- [5] J. Fisch, M. Henneaux, J. Stasheff, C. Teitelboim, *Existence, Uniqueness and Cohomology of the Classical BRST Charge with Ghosts of Ghosts*, Comm. Math. Phys. **120** (1989), 379-.
- [6] I.B. Frenkel, H. Garland, G.J. Zuckerman, *Semi-Infinite Cohomology and String Theory*, Proc. Natl. Acad. Sci. USA **83** (1987), 8442-8446.
- [7] J.F. Glazebrook, F.W. Kamber, *Chiral Anomalies and Dirac Families in Riemannian Foliations*, (1991) preprint.
- [8] B. Kostant, S. Sternberg, *Symplectic Reduction, BRS Cohomology, and Infinite Dimensional Clifford Algebras*, Ann. Phys. **176** (1987), 49-113.
- [9] R. Schmid, *Local Cohomology in Gauge Theories BRST Transformations and Anomalies*, Differential Geometry and its Applications, (1992) to appear.
- [10] R. Schmid, *BRST Cohomology and Anomalies*, Geometrical and Algebraic Aspects of Nonlinear Field Theory, 159-172, S. De Filippo et al (eds.), North - Holland, 1989.

- [11] R. Schmid, *BRST Bicomplexes*, preprint.
- [12] J. Stasheff, *The De Rham Bar Construction as a Setting for the Zumino, Faddeev, etc. Descent Equations*, Proc. Symp. anomalies, Geometry, Topology, Argonne Natl. Lab. 1984. W.E. Bardeen, A. R. White (eds.), World Scientific, Singapore (1985), 220-233. [Related papers can be found in the same proceedings].