

GROUPS OF OUTER TYPE E_6 WITH TRIVIAL TITS ALGEBRAS

SKIP GARIBALDI AND HOLGER P. PETERSSON

ABSTRACT. In two 1966 papers, J. Tits gave a construction of exceptional Lie algebras (hence implicitly exceptional algebraic groups) and a classification of possible indexes of simple algebraic groups. For the special case of his construction that gives groups of type E_6 , we connect the two papers by answering the question: Given an Albert algebra A and a separable quadratic field extension K , what is the index of the resulting algebraic group?

This article links two 1966 papers by Jacques Tits—namely, [Ti 66a] and [Ti 66b]—concerning simple linear algebraic groups over a field k .

In the first paper, he gave a construction that takes a composition algebra K and a degree 3 Jordan algebra A and produces a simple algebraic group $G(A, K)$. The Killing-Cartan type of the resulting group is given by the famous magic square:

		dim A					
		1	3	6	9	15	27
dim K	1			A_1	A_2	C_3	F_4
	2			A_2	$A_2 \times A_2$	A_5	E_6
	4	A_1	$A_1 \times A_1 \times A_1$	C_3	A_5	D_6	E_7
	8	G_2	D_4	F_4	E_6	E_7	E_8

In the second paper, he listed the possible indexes of a simple linear algebraic group. Recall from [Ti 66b, 2.3] that the index is the Dynkin diagram embellished so as to describe a maximal k -split torus and the action of the absolute Galois group on the Dynkin diagram.

The question

$$(0.1) \quad \textit{What is the index of the group } G(A, K)?$$

gives a natural link between the two papers. One wants a translation table between the index of $G(A, K)$ and properties of A and K . The action of the absolute Galois group is easy to see, so the serious part of Question (0.1) concerns the maximal k -split tori in $G(A, K)$.

The most interesting entries in the magic square are those in the bottom row and rightmost column. Of these, an answer to Question (0.1) is known

Date: July 24, 2006.

2000 Mathematics Subject Classification. Primary 20G15; Secondary 17C40, 17B25.

only for the upper right and lower left corners—corresponding to the “tautological” constructions $A \mapsto \text{Aut}(A)$ and $K \mapsto \text{Aut}(K)$ respectively [J 71, pp. 97, 98]—and the D_4 and F_4 in the bottom row. See Example 3.28 for a discussion of the two F_4 entries.

In this note, we completely settle Question (0.1) in the “next” interesting case, i.e., for the E_6 in the rightmost column. In that case, A is an Albert algebra (see §3 below for background) and K is a quadratic étale algebra. The hard part of answering the question is treated by the following theorem. We write $f_3(A)$ and $f_5(A)$ for the cohomological mod-2 invariants of A and refer to 3.24 for their precise definition. We fix an arbitrary base field k .

0.2. Theorem. *The following are equivalent:*

- (1) *The group $G(A, K)$ is isotropic.*
- (2) *$k \times K$ is (isomorphic to) a subalgebra of A .*
- (3) *A is reduced and there exists a 2-Pfister bilinear form γ such that*

$$\gamma \cdot f_3(A) = f_5(A) \quad \text{and} \quad \gamma \cdot [K] = 0.$$

Conditions (1)–(3) are implied by:

- (4) *A has a nonzero nilpotent element.*

Furthermore, if A is split by K , then (1)–(3) are equivalent to (4).

Once one knows that $G(A, K)$ is isotropic, it is not difficult to determine the index of $G(A, K)$ —see Prop. 2.3 and 4.8—so we have completely settled Question (0.1) for this entry in the magic square.

When K is “split” (i.e., $K = k \times k$, equivalently, $G(A, K)$ has type 1E_6), the theorem is a triviality. Indeed, conditions (1) through (3) are equivalent to the statement “ A is reduced”. When K is split, we define the statement “ A is split by K ” to mean that A is split as an Albert k -algebra.

According to [Ti 66b, p. 59], an isotropic group of type 2E_6 has one of eight possible indexes. Of these, only five can be the index of $G(A, K)$ for some Albert algebra A and quadratic étale algebra K . Conversely, every group with one of these five indexes is isogenous to $G(A, K)$ for some A and K , see Prop. 11.1. Roughly speaking, the various indexes are distinguished by their Rost invariant, see 2.8 and Prop. 2.9.

The main theorem shows the flavor of the paper; it mixes algebraic groups (in (1)), Jordan algebras (in (2) and (4)), and—essentially—quadratic forms (in (3)). The core of our proof is Jordan-theoretic. We prove (1) implies (2) or (4) in Cor. 5.3 and Propositions 7.2 and 8.1. We prove that (4) implies (2) in 6.3, (2) implies (3) in 6.2, and (3) implies (1) in 9.7. The last claim is proved in Example 5.4 and 10.1.

As side benefits of the proof, we obtain concrete descriptions of the projective homogeneous varieties for groups of type 2E_6 in §5 and we easily settle an open question from a 1969 paper of Veldkamp [V 69] in 5.5.

We remark that Tits’s construction paper excludes characteristics 2 and 3, but we extend his construction to work in all characteristics, see 4.4 and 4.5.

CONTENTS

1. Notation and reminders	3
2. Rost invariants	4
3. Albert algebra reminders	9
4. The groups $G(A, K)$	14
5. Homogeneous projective varieties	17
6. Embeddings of $k \times K$	20
7. The case where $T(x) = 0$	23
8. The case where $T(x) \neq 0$	26
9. Sufficient condition for isomorphism	27
10. The case where A is split by K	29
11. Applications to isotropic groups	29
12. Final observations	31
References	32

1. NOTATION AND REMINDERS

The *Tits algebras* of an algebraic group G are the k -algebras $\text{End}_G(V)$ as V varies over k -irreducible representations of G . We say that G has *trivial Tits algebras* if $\text{End}_G(V)$ is a (commutative) field for every V .

Below, K will always denote a quadratic étale k -algebra with nontrivial k -automorphism ι , and A is an Albert k -algebra.

Throughout, we use the notation $\langle \alpha_1, \dots, \alpha_n \rangle$ for the diagonal matrix with α_i in the (i, i) entry and for the symmetric bilinear form with that Gram matrix. We write $\langle\langle \alpha_1, \dots, \alpha_n \rangle\rangle$ for the Pfister bilinear from $\langle 1, -\alpha_1 \rangle \otimes \dots \otimes \langle 1, -\alpha_n \rangle$. *Pfister form* means ‘‘Pfister quadratic form’’. We write $I^n k$ for the module of quadratic forms generated by the n -Pfister forms over the Witt ring of symmetric bilinear forms (this agrees with the usual notation in characteristic $\neq 2$). We write $[K]$ for the 1-Pfister form given by the norm $K \rightarrow k$; a similar convention applies to the norm of an octonion k -algebra.

For $n \in \mathbb{N}$, we write $H^q(k, n)$ for the group denoted by $H^q(k, \mathbb{Z}/n\mathbb{Z}(q-1))$ in [GMS, pp. 151–155]. When n is not divisible by the characteristic of k , it is the Galois cohomology group $H^q(k, \mu_n^{\otimes(q-1)})$. There is a bijection between q -fold Pfister forms (up to isomorphism) and symbols in $H^q(k, 2)$. (The term ‘‘symbol’’ is defined to include the zero element of $H^q(k, 2)$. It corresponds to the hyperbolic q -Pfister form.) In characteristic different from 2, this is a direct consequence of Voevodsky’s proof of the Milnor Conjecture, and the bijection identifies the Pfister form $\langle\langle \alpha_1, \alpha_2, \dots, \alpha_q \rangle\rangle$ with the cup product $(\alpha_1) \cdot (\alpha_2) \cdots (\alpha_q)$. In characteristic 2 the bijection is described in [AB]. We will write, for example, $[K]$ also for the symbol in $H^1(k, 2)$ corresponding to the norm $K \rightarrow k$, and similarly for an octonion k -algebra.

2. ROST INVARIANTS

Let G be a quasi-simple, simply connected group over a field k . There is a canonical map $r_G: H^1(k, G) \rightarrow H^3(k, n_G)$ known as the *Rost invariant*, where n_G is a natural number depending on G , see [GMS] for details. The map is “functorial in k ”. In this section, we relate the Rost invariant with the index of isotropic groups of type 2E_6 with trivial Tits algebras.

2.1. There is a class $\nu \in H^1(k, \text{Aut}(G)^\circ)$ such that the twisted group G_ν is quasi-split. Moreover, this property uniquely determines ν , as can be seen from a twisting argument and the fact that the kernel of the map $H^1(k, \text{Aut}(G_\nu)^\circ) \rightarrow H^1(k, \text{Aut}(G_\nu))$ is zero.

If G has trivial Tits algebras, then there is an $\eta \in H^1(k, G)$ that maps to ν and we write $a(G) \in H^3(k, n_G)$ for the Rost invariant $r_G(\eta)$.

Lemma. *The element $a(G)$ depends only on the isomorphism class of G .*

Proof. Suppose that $\eta, \eta' \in H^1(k, G)$ both map to ν ; we prove that $\eta = \eta'$. Write G_η for the group G twisted by η , and write Z for the scheme-theoretic center of G_η . The sequence

$$1 \longrightarrow Z \longrightarrow G_\eta \longrightarrow \text{Aut}(G_\eta)^\circ \longrightarrow 1$$

is exact, and we find a commutative diagram

$$\begin{array}{ccccc} H_{\text{fppf}}^1(k, Z) & \longrightarrow & H^1(k, G_\eta) & \longrightarrow & H^1(k, \text{Aut}(G_\eta)^\circ) \\ & & \cong \downarrow \tau_\eta & & \cong \downarrow \\ & & H^1(k, G) & \longrightarrow & H^1(k, \text{Aut}(G)^\circ), \end{array}$$

where the vertical arrows are twisting isomorphisms. (Here, H^1 denotes Galois cohomology and H_{fppf}^1 denotes cohomology relative to the fppf site as in [DG] or [Wat]. The two notions agree for smooth groups like G .) Starting with $\eta' \in H^1(k, G)$, we find ν in the lower right, hence the trivial class in the upper right. Therefore, $\tau_\eta^{-1}(\eta')$ is in the image of $H_{\text{fppf}}^1(k, Z)$. However, G_η is quasi-split (by the choice of η), so it contains a quasi-trivial maximal k -torus S . Since Z is contained in S and $H^1(k, S)$ is zero, the image of $H_{\text{fppf}}^1(k, Z)$ in $H^1(k, G_\eta)$ is zero. Therefore, $\tau_\eta^{-1}(\eta')$ is zero and η' equals η . \square

2.2. Examples. In the following examples, G always denotes a quasi-simple, simply connected group with trivial Tits algebras. We use many facts from quadratic form theory. In characteristic $\neq 2$, the standard reference is [Lam]; in characteristic 2, some results and references are collected in [HL, p. 14]. The bijection between 3-Pfister forms and symbols in $H^3(k, 2)$ mentioned in §1 extends to a group isomorphism

$$e_3: I^3k/I^4k \xrightarrow{\sim} H^3(k, 2)$$

called the *Arason invariant*.

Suppose that q is an 8-dimensional form in I^3k . Then q is similar to a 3-Pfister form (see [Lam, X.5.6] in characteristic $\neq 2$) and $e_3(q)$ is a symbol. Since the correspondence between Pfister forms and symbols is a bijection, $e_3(q)$ is zero if and only if q is hyperbolic, if and only if q is isotropic.

- (a) Let G be of type 1D_n for $n = 3$ or 4 , so that G is isomorphic to $\text{Spin}(q)$ for some $2n$ -dimensional quadratic form q in I^3k , n_G is 2, and the invariant $a(G)$ is the Arason invariant of q . (See [KMRT, p. 437] in characteristic $\neq 2$; in characteristic 2, the same arguments apply.)

In the case $n = 3$, the Arason-Pfister Hauptsatz implies that q is hyperbolic, so G is split and $a(G)$ is zero.

In the case $n = 4$, q is similar to a 3-Pfister form. It follows that $a(G)$ is zero if and only if G is split if and only if G is isotropic.

- (b) Let G be of type 2D_n for $n = 3$ or 4 , with associated separable quadratic extension K/k . The group G is isomorphic to $\text{Spin}(q)$ for q a $2n$ -dimensional quadratic form such that $q - [K]$ is in I^3k , n_G is 2, and the Arason invariant of $q - [K]$ is $a(G)$ [KMRT, p. 437].

We claim that $a(G)$ is a symbol in $H^3(k, 2)$. By the remark at the start of this example, it suffices to prove that $q - [K]$ is Witt-equivalent to an 8-dimensional form. In case $n = 3$, $q \perp \langle -1 \rangle [K]$ is an 8-dimensional form and the claim is proved. In case $n = 4$, $q \perp \langle -1 \rangle [K]$ is a 10-dimensional form in I^3k , hence it is isotropic [Ti90, 4.4.1(ii)]. That is, $q \perp \langle -1 \rangle [K]$ is Witt-equivalent to an 8-dimensional form and the claim is proved.

We next observe that $a(G)$ is *not* killed by K if and only if $n = 4$ and G is k -anisotropic. In view of part (a), the “only if” direction is clear, so we prove the “if” direction and assume that $n = 4$ and G is k -anisotropic. For sake of contradiction, suppose that G is not anisotropic over K , i.e., that G is split by K , that is, q is hyperbolic over K . By the preceding paragraph, $q \perp \langle -1 \rangle [K]$ is isomorphic to $\langle \alpha \rangle \gamma \perp \mathcal{H}$ for some $\alpha \in k^\times$ and 3-Pfister form γ , where \mathcal{H} denotes a hyperbolic plane. As $q \perp \langle -1 \rangle [K]$ is K -hyperbolic, γ is isomorphic to $\beta[K]$ for some 2-Pfister bilinear form β by [HL, 4.2(iii)] and [GH, 1.1(i)]. Since $\langle \alpha \rangle \gamma \perp [K]$ is Witt-equivalent to q , it is isotropic (hence isomorphic to $q \perp \mathcal{H}$). But γ is k -anisotropic, so it represents $-\alpha\lambda$ for some nonzero λ represented by $[K]$. The roundness of γ and $[K]$ gives that $\langle \alpha \rangle \gamma$ is isomorphic to $\langle -\lambda \rangle \gamma$, hence to $\langle -1 \rangle \beta[K]$. Thus, $q \perp \mathcal{H}$ contains the subform $\langle 1, -1 \rangle [K]$, and q has Witt index at least 2. This contradicts the hypothesis that G is k -anisotropic, completing the proof of the observation.

- (c) Let G be of type 2A_5 . Then n_G is 2 [GMS, p. 146]. We prove: *If $a(G)$ is a symbol, then G is isotropic.*

The group G is isomorphic to the special unitary group of a K/k -hermitian form h [KMRT, 26.9], where K/k is the separable quadratic extension over which G is of type 1A_5 . We write q_h for the associated 12-dimensional quadratic trace of h over k as defined in [Sch, §10.1]. Note that G is isotropic if and only if h is isotropic if and only if q_h is isotropic [Sch, 10.1.1].

Suppose that $a(G)$ is a symbol, i.e., $a(G)$ equals $e_3(\gamma)$ for some 3-Pfister form γ . The function field F of γ kills $a(G)$, so splits G (because the Rost invariant of the quasi-split 2A_5 has zero kernel [Ga, 1.10]), hence makes q_h hyperbolic. Therefore, the anisotropic part of q_h is of the form $\tau \otimes \gamma$ for some bilinear form τ . But 8 (the dimension of γ) does not divide 12 (the dimension of q_h), so we conclude that q_h is isotropic over k .

The purpose of the preceding examples was to prepare the proof of the following proposition.

2.3. Proposition. *Let G be an isotropic group of type 2E_6 with trivial Tits algebras; write K for the associated quadratic extension of k . Then $a(G)$ is in $H^3(k, 2)$ and the index of G is given by Table 2.4.*





index	condition
quasi-split	$a(G) = 0$
	$a(G)$ is a nonzero symbol killed by K
	$a(G)$ is a symbol not killed by K
	$a(G)$ is not a symbol

TABLE 2.4. Tits indexes and their corresponding Rost invariants

Proof. Consulting the list of possible indexes from [Ti66b, p. 59], the only one missing from the table is



But a group with such an index has nontrivial Tits algebras [Ti71, p. 211], therefore the index of G is in the table.

If G is not quasi-split, then the semisimple anisotropic kernel H of G is quasi-simple with trivial Tits algebras. It follows from Tits's Witt-type theorem that the class η from 2.1 is the image of some $\eta_H \in H^1(k, H)$,

chosen so that H twisted by η_H is quasi-split. Then $a(H)$ equals $r_H(\eta_H)$. The composition

$$H^1(k, H) \longrightarrow H^1(k, G) \xrightarrow{r_G} H^3(k, n_G)$$

equals r_H by [GMS, p. 122] because the inclusion $H \subset G$ comes from the obvious inclusion of root systems. Therefore,

$$a(H) = r_H(\eta_H) = r_G(\eta) = a(G).$$

The correspondence between the index and $a(G)$ asserted by the table now follows from Example 2.2. \square

2.5. Suppose that G is a quasi-simple, simply connected group with trivial Tits algebras, and that G' is a *strongly inner form* of G , that is, G' can be obtained by twisting G by a 1-cocycle $\alpha \in Z^1(k, G)$. A twisting argument combined with [Gi, p. 76, Lemme 7] gives: $a(G)$ equals $a(G')$ if and only if $r_G(\alpha)$ is zero.

Write Z for the scheme-theoretic center of G . The group $H_{\text{fppf}}^1(k, Z)$ acts on $H^1(k, G)$, and the same twisting argument shows that $r_G(\alpha)$ equals $r_G(\zeta \cdot \alpha)$ for all $\zeta \in H_{\text{fppf}}^1(k, Z)$ and $\alpha \in H^1(k, G)$.

Sometimes a group G is determined up to isomorphism by $a(G)$ and a few additional data.

2.6. Lemma. *Let G be a simply connected group of type 1D_4 or 2D_4 with trivial Tits algebras, and let G' be a strongly inner form of G . If $a(G)$ equals $a(G')$, then G and G' are isomorphic.*

Proof. We identify G' with the group G_α obtained by twisting G by some 1-cocycle α , and we assume that $r_G(\alpha)$ is zero. We identify G with $\text{Spin}(q)$ for some 8-dimensional quadratic form q , which identifies G' with $\text{Spin}(q_\alpha)$ such that

$$(2.7) \quad 0 = r_G(\alpha) = e_3(q_\alpha - q).$$


However, G_α is the same as $G_{\zeta \cdot \alpha}$ for all $\zeta \in H_{\text{fppf}}^1(k, Z)$, and (2.7) holds also when α is replaced by $\zeta \cdot \alpha$. In particular, $q_{\zeta \cdot \alpha} - q$ belongs to $I^4 k$ for all ζ .

The vector representation of G restricts to a surjection $\chi: Z \rightarrow \mu_2$, hence to a map

$$\chi: H_{\text{fppf}}^1(k, Z) \rightarrow H_{\text{fppf}}^1(k, \mu_2) = k^\times / k^{\times 2},$$

and the form $q_{\zeta \cdot \alpha}$ is isomorphic to $\langle \chi(\zeta) \rangle q_\alpha$. Choose ζ so that $q_{\zeta \cdot \alpha}$ and q represent a common element in k^\times . Then $q_{\zeta \cdot \alpha} - q$ is isotropic, hence has anisotropic part of dimension ≤ 14 . However, that form also belongs to $I^4 k$, so it is hyperbolic by the Arason-Pfister Hauptsatz. We conclude that $q_{\zeta \cdot \alpha}$ is isomorphic to q_α , hence G' is isomorphic to G . \square

2.8. Let G be a simply connected group of type E_6 that is split or has index



In either case, G has trivial Tits algebras [Ti 71, 6.4.5]. The Rost invariant of G has zero kernel by [GH, 5.5], so by 2.5 we have: *If G' is a strongly inner form of G such that $a(G') = a(G)$, then G' is isomorphic to G .*

The same argument as in the proof of Prop. 2.3 shows that the index of G is determined by $a(G)$: Obviously $a(G)$ is zero if G is split, and if G has the displayed index, then $a(G)$ is a nonzero symbol in $H^3(k, 2)$ by Example 2.2a.

A group G from the top three rows of Table 2.4 is completely determined by the value of $a(G)$.

2.9. Proposition. *Let G be a simply connected group of type 2E_6 , and let G' be a strongly inner form of G . Suppose further that the indexes of G and G' appear in the top three lines of Table 2.4. If $a(G)$ equals $a(G')$, then G and G' are isomorphic.*

Example 11.4 below shows that the hypothesis on the indexes of G and G' is necessary.

Proof. Fix a maximal k -split torus S in G , a maximal k -torus T containing it, and a set of simple roots for G with respect to T . Since G is simply connected, the group of cocharacters of T is identified with the coroot lattice. Write S_1 for the rank 1 torus corresponding to the sum of fundamental weights $\omega_1 + \omega_6$ (“corresponding to the circle around the α_1 and α_6 vertices in the index”); it is k -defined by [BT, Cor. 6.9]. Put G_1 for the derived subgroup of $Z_G(S_1)$; it is simply connected of type 2D_4 with trivial Tits algebras; it has the same semisimple anisotropic kernel as G .

Define a subgroup G'_1 of G' in an analogous manner. Since G_1 and G'_1 are both of type 2D_4 with trivial Tits algebras, G'_1 is isomorphic to G_1 twisted by some 1-cocycle $\alpha \in Z^1(k, G_1)$.

As in the proof of Prop. 2.3, $r_{G_1}(\alpha)$ equals $r_G(\alpha)$, which is zero by 2.5. Lemma 2.6 shows that G'_1 is isomorphic to G_1 , hence the semisimple anisotropic kernels of G and G' agree. Tits’s Witt-type theorem gives that G and G' are isomorphic. \square

2.10. Remark ($\text{char } k \neq 2$). Let G be as in Prop. 2.9 and let γ be the 3-Pfister form corresponding to $a(G)$. We claim that the group G_1 in the proof of the proposition is isomorphic to $\text{Spin}(\gamma^K)$, where γ^K denotes the K -associate of γ as defined in [KMRT, p. 499]. In particular, *the semisimple anisotropic kernel of G is the semisimple anisotropic kernel of $\text{Spin}(\gamma^K)$* . To prove the claim, let G' be the quasi-split strongly inner form of G , so G'_1 is the spin group of a quasi-split quadratic form q . Fix $\alpha \in H^1(k, \text{Spin}(q))$ such that q_α is isomorphic to γ^K . The twisted group G'_α is as in the first three lines of the table and $a(G'_\alpha)$ equals the Arason invariant of $q_\alpha - q$. This form is Witt-equivalent to $\langle \delta \rangle \gamma$ for $\delta \in k^\times$ such that $K = k(\sqrt{\delta})$, hence $a(G'_\alpha)$ equals $a(G)$ and G'_α is isomorphic to G by Prop. 2.9. Since the semisimple anisotropic kernels of G'_α and $\text{Spin}(q_\alpha)$ agree, the claim follows.

3. ALBERT ALGEBRA REMINDERS

3.1. Arbitrary Albert algebras. Albert algebras are Jordan algebras of degree 3 and hence may all be obtained from cubic forms with adjoint and base point in the sense of [McC 69]. More specifically, given an Albert algebra A over k , there exist a cubic form $N : A \rightarrow k$ (the *norm*) and a quadratic map $\sharp : A \rightarrow A, x \mapsto x^\sharp$ (the *adjoint*) which, together with the unit element $1 \in A$, satisfy the relations $N(1) = 1, 1^\sharp = 1$,

$$(3.2) \quad \begin{aligned} x^\sharp &= N(x)x, \\ (DN)(x)y &= T(x^\sharp, y), \end{aligned}$$

$$(3.3) \quad 1 \times x = T(x)1 - x$$

in all scalar extensions, where $T = -(D^2 \log N)(1) : A \times A \rightarrow k$ (the *trace form*) stands for the logarithmic hessian of N at 1, $T(x) = T(x, 1)$ and $x \times y = (x + y)^\sharp - x^\sharp - y^\sharp$ is the bilinearization of the adjoint. The U -operator of A is then given by the formula

$$(3.4) \quad U_x y = T(x, y)x - x^\sharp \times y.$$

The *quadratic trace*, defined by $S(x) := T(x^\sharp)$, is a quadratic form with bilinearization

$$(3.5) \quad S(x, y) = T(x)T(y) - T(x, y).$$

It relates to the adjoint by the formula

$$(3.6) \quad x^\sharp = x^2 - T(x)x + S(x)1,$$

where, as in [J68, 1.5], the powers of $x \in A$ are defined by $x^0 = 1, x^1 = x$, and $x^{n+2} = U_x x^n$ for $n \geq 0$. For future use, we recall the following identities from [McC 69].

$$(3.7) \quad x^\sharp \times (x \times y) = N(x)y + T(x^\sharp, y)x,$$

$$(3.8) \quad T(x^\sharp, x) = 3N(x),$$

$$(3.9) \quad x^\sharp \times x = [S(x)T(x) - N(x)]1 - S(x)x - T(x)x^\sharp,$$

and

$$(3.10) \quad x^\sharp \times (y \times z) + (x \times y) \times (x \times z) = T(x^\sharp, y)z + T(x^\sharp, z)y + T(y \times z, x)x.$$

Just as in general Jordan rings, the *Jordan triple product* derives from the U -operator (3.4) through linearization:

$$(3.11) \quad \begin{aligned} \{x, y, z\} &:= U_{x,z}y := (U_{x+y} - U_x - U_y)z \\ &= T(x, y)z + T(y, z)x - (z \times x) \times y. \end{aligned}$$

Finally, the generic minimum polynomial of x in the sense of [JK] is

$$(3.12) \quad m_x(\mathbf{t}) = \mathbf{t}^3 - T(x)\mathbf{t}^2 + S(x)\mathbf{t} - N(x) \in k[\mathbf{t}],$$

so by [JK, p. 219] we have

(3.13)

$$x^3 - T(x)x^2 + S(x)x - N(x)1 = 0 = x^4 - T(x)x^3 + S(x)x^2 - N(x)x,$$

where the second equation is a *trivial* consequence of the first only for char $k \neq 2$ because then we are dealing with *linear* Jordan algebras. An element $x \in A$ is invertible if and only if $N(x) \neq 0$. At the other extreme, $x \in A$ is said to be *singular* if $x^\sharp = 0 \neq x$. As an ad-hoc definition, singular idempotents will be called *primitive*.

The following lemma collects well-known properties of the Peirce decomposition relative to primitive idempotents, using the labelling of [Lo 75]. We refer to [Fa, Lemma 1.5], [Ra 72, (28), (31), Lemma 2], and [PR 95, Lemma 5.3d] for details.

3.14. Lemma. *Let $e \in A$ be a primitive idempotent and put $f = 1 - e$.*

(a) *The Peirce components of A relative to e are described by the relations*

$$\begin{aligned} A_2(e) &= ke, \\ A_1(e) &= \{x \in A \mid T(x) = 0, e \times x = 0\}, \\ A_0(e) &= \{x \in A \mid e \times x = T(x)f - x\}. \end{aligned}$$

(b) *The restriction S_e of S to $A_0(e)$ is a quadratic form with base point f over k whose associated Jordan algebra agrees with $A_0(e)$ as a subalgebra of A .*

(c) *$x^\sharp = S(x)e$ for all $x \in A_0(e)$.* □

3.15. Reduced Albert algebras. An Albert k -algebra A is *reduced* if it is not division, i.e., if it contains nonzero elements that are not invertible. Every such algebra is isomorphic to a Jordan algebra $\mathcal{H}_3(C, \Gamma)$ as defined in the following paragraph.

Let C be an octonion (or Cayley) k -algebra (see [KMRT, §33.C] for the definition and elementary properties) and fix a diagonal matrix $\Gamma := \langle \gamma_1, \gamma_2, \gamma_3 \rangle \in \text{GL}_3(k)$. We write $\mathcal{H}_3(C, \Gamma)$ for the vector space of 3-by-3 matrices x with entries in C that are Γ -hermitian ($x = \Gamma^{-1}\bar{x}^t\Gamma$) and have scalars down the diagonal. It is spanned by the diagonal unit vectors e_{ii} ($1 \leq i \leq 3$) and by the hermitian matrix units $x_i[jl] := \gamma_l x_i e_{jl} + \gamma_j \bar{x}_i e_{lj}$ ($x_i \in C$), where (ijl) here and in the sequel always varies over the cyclic permutations of (123). It has a Jordan algebra structure derived from a cubic form with adjoint and base point as in 3.1: Writing N_C for the norm, T_C for the trace and $x \mapsto \bar{x}$ for the conjugation of C , we let

$$x = \sum \alpha_i e_{ii} + \sum x_i[jl], \quad y = \sum \beta_i e_{ii} + \sum y_i[jl], \quad (\alpha_i, \beta_i \in k, x_i, y_i \in C)$$

be arbitrary elements of A and set

$$(3.16) \quad N(x) = \alpha_1 \alpha_2 \alpha_3 - \sum \gamma_j \gamma_l \alpha_i N_C(x_i) + \gamma_1 \gamma_2 \gamma_3 T_C(x_1 x_2 x_3),$$

$$(3.17) \quad x^\sharp = \sum (\alpha_j \alpha_l - \gamma_j \gamma_l N_C(x_i)) e_{ii} + \sum (\gamma_i \bar{x}_j \bar{x}_l - \alpha_i x_i) [jl],$$

as well as $1 = \sum e_{ii}$. Then $(N, \sharp, 1)$ is a cubic form with adjoint and base point whose associated trace form is

$$(3.18) \quad T(x, y) = \sum \alpha_i \beta_i + \sum \gamma_j \gamma_l N_C(x_i, y_i).$$

The Jordan algebra structure on $\mathcal{H}_3(C, \Gamma)$ is defined to be the one associated with $(N, \sharp, 1)$.

Specializing y to 1 in (3.18) yields

$$(3.19) \quad T(x) = \sum \alpha_i,$$

whereas linearizing (3.17) leads to the relation

$$(3.20) \quad x \times y = \sum (\alpha_j \beta_l + \beta_j \alpha_l - \gamma_j \gamma_l N_C(x_i, y_i)) e_{ii} \\ + \sum (\gamma_i \overline{x_j y_l} + \overline{y_j x_l} - \alpha_i y_i - \beta_i x_i) [jl].$$

Furthermore, the quadratic trace S of A and its polarization by (3.17), (3.19), (3.20) have the form

$$(3.21) \quad S(x) = \sum (\alpha_j \alpha_l - \gamma_j \gamma_l N_C(x_i)), \\ S(x, y) = \sum (\alpha_j \beta_l + \beta_j \alpha_l - \gamma_j \gamma_l N_C(x_i, y_i)).$$

The triple $F := (e_{11}, e_{22}, e_{33})$ is a *frame*, i.e., a complete orthogonal system of primitive idempotents, in A with Peirce components $J_{ii}(F) = k e_{ii}$, $J_{jl}(F) = C[jl]$. Comparing this with (3.21), we obtain a natural isometry

$$(3.22) \quad S|_{J_{jl}(F)} \cong \langle -\gamma_j \gamma_l \rangle \cdot N_C.$$

3.23. The split Albert algebra. Of particular interest is the *split* Albert algebra $A^d = \mathcal{H}_3(C^d, 1)$, where C^d stands for the split octonion algebra of Zorn vector matrices over k [KMRT, Ch. VIII, Exercise 5] and 1 is the 3-by-3 unit matrix. Albert algebras are exactly the k -forms of A^d , i.e., they become isomorphic to A^d over the separable closure of k . For any Albert algebra A over k as in 3.1, the equation $N = 0$ defines a singular hypersurface in $\mathbb{P}(A)$, its singular locus being given by the system of quadratic equations $x^\sharp = 0$ in $\mathbb{P}(A)$.

3.24. The invariants f_3 and f_5 . Letting A be a reduced Albert algebra over k as in 3.15, we briefly describe the cohomological mod-2 invariants of A in a characteristic-free manner; for $\text{char } k \neq 2$, see [GMS, pp. 49–52]. Since C is uniquely determined by A up to isomorphism [Fa, Th. 1.8], so is its norm N_C , which, being a 3-fold Pfister form, gives rise to the first mod-2 invariant $f_3(A) \in H^3(k, 2)$ of A . On the other hand, Γ is not uniquely determined by A since, for example, it may be multiplied by nonzero scalars and its components may be permuted arbitrarily as well as multiplied by nonzero square factors without changing the isomorphism class of A . This

allows us to assume $\gamma_2 = 1$. Then, as in [P, 2.1, 4.1], we may consider the 5-fold Pfister form $\langle\langle -\gamma_1, -\gamma_3 \rangle\rangle N_C$, giving rise to the second mod-2 invariant

$$(3.25) \quad f_5(A) = f_3(A) \cdot (-\gamma_1) \cdot (-\gamma_3) \in H^5(k, 2)$$

of A . Translating Racine's characteristic-free version [Ra 72, Th. 3] of Springer's classical result [J 68, p. 381, Th. 6] to the cohomological setting, it follows in all characteristics that reduced Albert algebras are classified by their mod-2 invariants f_3, f_5 .

3.26. Nilpotent elements. Let A be an Albert algebra over k . An element $x \in A$ is said to be *nilpotent* if $x^n = 0$ for some $n \in \mathbb{N}$. Combining [JK, p. 222, Th. 2(vi)] with (3.12), we conclude that

$$(3.27) \quad x \in A \text{ is nilpotent if and only if } T(x) = S(x) = N(x) = 0.$$

In this case $x^3 = x^4 = 0$ by (3.13). Hence A contains nonzero nilpotents if and only if $x^2 = 0$ for some nonzero element $x \in A$. We also conclude from (3.6), (3.27) that $x \in A$ satisfies $x^2 = 0$ if and only if $x^\sharp = 0$ and $T(x) = 0$. Finally, by [P, 4.4], an Albert algebra contains nonzero nilpotent elements if and only if it is reduced and $f_5(A) = 0$.

3.28. Example. ($\text{char } k \neq 2, 3$) We recall the solution to Question (0.1) for the two F_4 entries. The one in the upper right corner of the magic square amounts to the map $A \mapsto \text{Aut}(A)$. There are three possible indexes for $\text{Aut}(A)$, corresponding to the cases where $\text{Aut}(A)$ is split, $\text{Aut}(A)$ is isotropic but not split, and $\text{Aut}(A)$ is anisotropic. It follows from the description of the projective homogeneous varieties of $\text{Aut}(A)$ —see e.g. [CG, 9.1, 9.4]—that $\text{Aut}(A)$ is split if and only if A is the split Albert algebra and that $\text{Aut}(A)$ is isotropic if and only if A contains a nonzero nilpotent. This gives an answer to Question (0.1) that is similar to our Cor. 5.3 below. A more satisfying answer is in terms of the cohomological invariants of A : a criteria for nilpotents was given in the last sentence of 3.26, and the algebra A is split if and only if it is reduced (detected by Rost's mod 3 invariant) and $f_3(A)$ is zero.

As for the other F_4 entry in the magic square, one knows that every central simple Jordan algebra of degree 3 and dimension 6 is of the form $\mathcal{H}_3(k, \Gamma)$ for a symmetric $\Gamma \in GL_3(k)$. Further, Tits's group of type F_4 constructed from it and an octonion algebra C is isomorphic to the automorphism group of the reduced Albert algebra $\mathcal{H}_3(C, \Gamma)$ by [Wal, p. 51]. Therefore, the answer to Question (0.1) for this entry is the same as for the previous entry.

In order to bring Jordan-theoretic techniques to bear in the proof of Theorem 0.2, we will translate hypothesis (1) into a condition on the Albert algebra A . To do so, we extend the results of [CG, §13] to the case where k is arbitrary.

3.29. Definition. A subspace X in an Albert algebra A is an *inner ideal* if $U_x A \subseteq X$ for all $x \in X$. A subspace X of A is *singular* if $x^\sharp = 0$ for all

$x \in X$. A subspace X of A is a *hyperline* if it is of the form $x \times A$ for some nonzero x with $x^\sharp = 0$.

The nonzero, proper inner ideals of an Albert algebra A are the singular subspaces (of dimensions 1 through 6) and the hyperlines (of dimension 10), see [McC 71, p. 467] and [Ra 77, Th. 2].

For an inner ideal X of A , we define $\psi(X)$ to be the set of all $y \in A$ satisfying

$$(3.30) \quad \{X, y, A\} \subseteq X$$

and

$$(3.31) \quad U_A U_y X \subseteq X.$$

The definition of ψ in [CG, 13.1] consisted of only condition (3.30). Condition (3.31) was suggested by Erhard Neher, for the purposes of including characteristic 2. He points out that when k has characteristic not 2, (3.31) follows from (3.30) by applying the identity JP13. (The notation JP xx refers to the Jordan pair identities as numbered in [Lo 75].)

3.32. Lemma (Neher). *If X is an inner ideal of A , then $\psi(X)$ is also an inner ideal.*

Proof. Clearly $\psi(X)$ is closed under scalar multiplication. Let y, z be elements of $\psi(X)$. To prove that $y + z$ is in $\psi(X)$, it suffices to show that $U_A U_{y,z} X \subseteq X$, which follows from JP13:

$$U_a U_{y,z} x = \{a, y, \{a, z, x\}\} - \{U_a y, z, x\} \in X.$$

Fix $y \in \psi(X)$ and $a \in A$. By JP7, we have

$$\{X, U_y a, A\} \subseteq \{\{a, y, X\}, y, A\} + \{a, U_y X, A\} \subseteq X,$$

So $U_y a$ satisfies condition (3.30). Also, by JP3,

$$U_A U_{U_y a} X = U_A U_y U_a U_y X \subseteq X,$$

so $U_y a$ satisfies condition (3.31). \square

We now check that the descriptions of $\psi(X)$ for various X given in [CG, §13] are valid in all characteristics. Consider first the case where X is 1-dimensional singular, i.e., an “ α_1 -space” in the language of [CG]. For $y \in X \times A$, we have $\{X, y, A\} \subseteq X$ by [CG, 13.6]. The argument in the last paragraph of the proof of [CG, Prop. 13.19] shows that $U_y X$ is zero. Therefore, $\psi(X)$ contains $X \times A$, and as in [CG, 13.8]—using that $\psi(X)$ is an inner ideal—we conclude that $\psi(X)$ equals $X \times A$.

Similar arguments easily extend the proofs of 13.9–13.19 of [CG] to include the case where k has arbitrary characteristic. (See also Remarks 5.5 and 13.22 of that paper.) For every x in a proper inner ideal X and every $y \in \psi(X)$, we find *a posteriori* that $U_y x$ is zero. That is, (3.31) is superfluous.

3.33. In summary, for a proper, nonzero inner ideal X in A , the set $\psi(X)$ of $y \in A$ satisfying (3.30) is an inner ideal of A whose dimension is given by

the following table. The 5-dimensional inner ideals come in two flavors; we write “5’” to indicate a 5-dimensional maximal singular subspace.

$\dim X$	1	2	3	5'	6	10
$\dim \psi(X)$	10	5'	3	2	6	1

For X of dimension $\neq 6$, the space $\psi(X)$ is the set of all $y \in A$ such that $\{X, y, A\}$ is zero. For X of dimension 6, $\psi(X)$ is a 6-dimensional inner ideal such that $\{X, \psi(X), A\}$ equals X .

4. THE GROUPS $G(A, K)$

This section defines the groups $G(A, K)$ using *hermitian Jordan triples*. These triples were previously studied in an analytic setting [Lo 77, 2.9]; translating to a purely algebraic situation in a natural way, we arrive at the following concept.

4.1. Definition. A *hermitian Jordan triple* over k is a triple (K, V, P) consisting of a quadratic étale k -algebra K (with conjugation ι), a free K -module V of finite rank, and a quadratic map $P: V \rightarrow \text{Hom}_K(V^\iota, V)$, where V^ι is the K -module V with scalar multiplication twisted by ι , such that (V, P) is an ordinary Jordan triple over k in the sense of, e.g., [Lo 75, 1.13]. In particular, $P_v: V \rightarrow V$ is an ι -semilinear map depending K -quadratically on $v \in V$.

A homomorphism $(K, V, P) \rightarrow (K', V', P')$ of hermitian Jordan triples is a pair (ϕ, h) such that $\phi: K \xrightarrow{\sim} K'$ is a k -isomorphism and $h: V \rightarrow V'$ is a ϕ -semilinear map such that $h(P_x y) = P_{h(x)} h(y)$ for all $x, y \in V$.

It is easy to see that Jordan pairs are basically the same as hermitian Jordan triples (K, V, P) with K split.

4.2. Example. Starting with a Jordan k -algebra J and a quadratic étale k -algebra K with conjugation ι , we obtain a hermitian Jordan triple as output. Namely, take $V = J \otimes_k K$ and define P by

$$P_x y := U_x (\text{Id}_J \otimes \iota) y.$$

We denote this hermitian Jordan triple by $\mathcal{T}(J, K)$.

A *hermitian Albert triple* is a triple $\mathcal{T}(A, K)$, where A is an Albert k -algebra. We call the triple $\mathcal{T}^d := \mathcal{T}(A^d, k \times k)$ the *split Albert triple* or simply the *split triple*.

We will now explicitly describe the automorphism group of a hermitian Jordan triple $\mathcal{T}(J, K)$. Following [J 81, 1.7, p. 1.23], the *structure group* of J , denoted by $\text{Str}(J)$, is the subgroup of $\text{GL}(J)$ consisting of all bijective linear maps $g: J \rightarrow J$ that may be viewed as isomorphisms from J onto an appropriate isotope or, equivalently, that have the property that there exists a bijective linear map $g^\dagger: J \rightarrow J$ satisfying the relation $U_{g(x)} = g U_x (g^\dagger)^{-1}$ for all $x \in J$; obviously it is a group scheme. The assignment $g \mapsto g^\dagger$ is an order 2 automorphism of the structure group of J .

4.3. Lemma. *The group of k -automorphisms of $\mathcal{T}(J, K)$ is generated by the element $(\iota, \text{Id}_J \otimes \iota)$ of order 2 and by the elements (Id_K, g) for $g \in \text{Str}(J)(K)$ satisfying $g^\dagger = (\text{Id}_J \otimes \iota)g(\text{Id}_J \otimes \iota)$.*

Proof. $(\iota, \text{Id}_J \otimes \iota)$ is clearly an automorphism of $\mathcal{T}(J, K)$. Conversely, multiplying any automorphism (ϕ, h) of $\mathcal{T}(J, K)$ by $(\iota, \text{Id}_J \otimes \iota)$ if necessary, we are allowed to assume that ϕ is the identity on K . The equation $h(P_x y) = P_{h(x)} h(y)$ is equivalent to $U_{h(x)} = hU_x(\text{Id}_J \otimes \iota)h^{-1}(\text{Id}_J \otimes \iota)$, which gives the claim. \square

4.4. Since the trace form of an Albert algebra A is a non-degenerate symmetric bilinear form, it follows from [McC69, p. 502] that the structure group of A agrees with its group of norm similarities; viewed as a group scheme, it is a reductive algebraic group with center of rank 1 and its semisimple part is simply connected of type E_6 [Spr, 14.2].

Below, we only consider hermitian Jordan triples $\mathcal{T} = (V, K, P)$ that become isomorphic to \mathcal{T}^d over a separable closure k_{sep} of k . We write $G(\mathcal{T})$ for the semisimple part of the identity component of $\text{Aut}(\mathcal{T})$; when \mathcal{T} is of the form $\mathcal{T}(A, K)$, we will write $G(A, K)$ for short. This group is simply connected of type E_6 because it is isomorphic to the semisimple part of $\text{Str}(A^d)$ over k_{sep} . It acts on the vector space underlying \mathcal{T} , and over a separable closure this representation is isomorphic to a direct sum of the usual representation of E_6 on A^d and its contragredient. Since the direct sum of these two representations is defined over k , the group has trivial Tits algebras. Also, $G(\mathcal{T})$ is of type 2E_6 if and only if K is a field.

4.5. Relation with Tits's construction. We now observe that the Lie algebra \mathfrak{g} of $G(A, K)$ is precisely the one obtained from A and K using Tits's construction from [Ti66a]. For the duration of this subsection, we assume — as in [Ti66a] — that the characteristic of k is $\neq 2, 3$. Suppose first that K is $k \times k$. From the first paragraph of 4.4, we see that \mathfrak{g} consists of those linear transformations on A that leave the cubic form N “Lie invariant”. By [J59, p. 189, Th. 5], this is the same as the Lie algebra $R_0(A) \oplus D(A)$, where $R_0(A)$ is the vector space generated by the transformations “right multiplication by a trace zero element of A ” and $D(A)$ denotes the derivation algebra of A ; this is what one obtains from Tits's construction.

Now consider the case where K is a field. We may view the vector space $A \otimes K$ underlying $\mathcal{T}(A, K)$ as the subspace of $A_K \times A_K$ fixed by the map $(x, y) \mapsto (\iota y, \iota x)$. In this way, we can view $G(A, K)$ as the group whose k -points are those $(f, f^\dagger) \in G(A_K, K \times K)(K)$ such that $\iota f^\dagger \iota = f$. The differential of $f \mapsto f^\dagger$ is $d \mapsto -d^*$. It follows that the Lie algebra \mathfrak{g} is the subalgebra of $R_0(A_K) \oplus D(A_K)$ consisting of elements fixed by the map $d \mapsto -\iota d^* \iota$. When d is right multiplication by an element of A_K , we have $d^* = d$ because the bilinear form T is associative [J68, p. 227, Cor. 4]. The derivation algebra $D(A_K)$ is spanned by commutators of right multiplications, hence is fixed elementwise by the map $d \mapsto -d^*$. Consequently, the

Lie algebra of $G(A, K)$ is identified with $\sqrt{\lambda}R_0(A) \oplus D(A)$, where $\lambda \in k^\times$ satisfies $K = k(\sqrt{\lambda})$. This is the Lie algebra constructed by Tits. (Jacobson [J 71] and Ferrar [Fe 69] denote this Lie algebra by $\mathcal{E}_6(A)_\lambda$ and $\mathcal{L}(A)_\lambda$ respectively.)

By Galois descent, the map $\mathcal{T} \mapsto G(\mathcal{T})$ produces (up to isogeny) all groups of type E_6 with trivial Tits algebras over every field k . But over some fields, it suffices to consider only the triples of the form $\mathcal{T}(A, K)$.

4.6. Example. Every simply connected group of type 1E_6 with trivial Tits algebras can be realized as the group of linear transformations on an Albert algebra A preserving the cubic norm, hence is of the form $G(A, k \times k)$. (In characteristic $\neq 2, 3$, this follows from [Ga, 3.4].) We now list a few examples of fields k such that every simply connected group of type 2E_6 with trivial Tits algebras is—up to isogeny—of the form $G(A, K)$.

- (a) When k is a local field [Kne] or a finite field, every group of type 2E_6 is quasi-split.
- (b) For k the real numbers, up to isogeny the three groups of type 2E_6 are of the form $G(A, \mathbb{C})$ as A varies over the three isomorphism classes of Albert \mathbb{R} -algebras [J 71, pp. 119, 120].
- (c) When k is a number field, as can be seen by using the Hasse Principle, cf. [Fe 78, 3.2, 6.4].

For f an automorphism of an Albert algebra A , we have $f^\dagger = f$, so $\text{Aut}(A)$ is a subgroup of $G(A, K)$ for all K .

4.7. Example. The group $G(A^d, K)$ contains the split group $\text{Aut}(A^d)$ of type F_4 , hence has k -rank at least 4. The classification of possible indexes gives that $G(A^d, K)$ is quasi-split for all K , including the case $K = k \times k$.

4.8. Rost invariant of $G(A, K)$. The inclusion of $\text{Aut}(A)$ in $G(A, K)$ has Rost multiplier one [Dy, p. 194] in the language of [GMS], hence the composition

$$H^1(k, \text{Aut}(A)) \longrightarrow H^1(k, G(A, K)) \xrightarrow{r_{G(A, K)}} H^3(k, n_{G(A, K)})$$

agrees with the Rost invariant relative to $\text{Aut}(A)$. In the definition of $a(G)$ at the start of §2, one can take the image of A^d as η by the preceding example. We find:

$$a(G(A, K)) = -r_{F_4}(A) \in H^3(k, 6),$$

where the right side denotes the negative of the usual Rost invariant of the Albert algebra A . In particular, the mod-2 portion of $a(G(A, K))$ is the symbol $f_3(A)$.

If one knows that a group $G(A, K)$ is isotropic, then combining the previous paragraph and Prop. 2.3 gives the index of $G(A, K)$.

5.2. Proposition. *The projective homogeneous variety for $G(\mathcal{T})$ corresponding to the subset S of the Dynkin diagram has k -points the inner ideals as in the table below.*

S	k -points are inner ideals
$\{\alpha_1, \alpha_6\}$	$X \subset Y$ s.t. $\text{rank } X = 1$, $\text{rank } Y = 10$, and $[X, Y, V] = 0$
$\{\alpha_3, \alpha_5\}$	$X \subset Y$ s.t. $\text{rank } X = 2$, $\text{rank } Y = 5$, and $[X, Y, V] = 0$
$\{\alpha_4\}$	X s.t. $\text{rank } X = 3$ and $[X, X, V] = 0$
$\{\alpha_2\}$	X s.t. $\text{rank } X = 6$ and $[X, X, V] = X$

With the information from the table, it should be no trouble to describe the homogeneous varieties corresponding to other opposition-stable subsets of the Dynkin diagram using the recipe in [CG, 9.4]. (The *opposition involution* is the unique nonidentity automorphism of the Dynkin diagram.)

Proof. Since the claimed collections of subspaces form projective k -varieties on which $G(\mathcal{T})$ acts, it remains only to show that the action is transitive on k_{sep} -points and that the stabilizer of a k_{sep} -point is a parabolic subgroup of the correct type. Therefore, we assume that k is separably closed—in particular that \mathcal{T} is split, i.e., isomorphic to \mathcal{T}^d —and note that $G(\mathcal{T})$ is the split simply connected group of type E_6 . The projective homogeneous variety Z_0 associated with the split group and S was described in sections 7 and 9 of [CG]. We prove the proposition by producing a $G(\mathcal{T})(k)$ -equivariant bijection $Z(k) \xrightarrow{\sim} Z_0(k)$.

Let $X \subset Y$ be inner ideals as in the first line of the table. As in Example 5.1, we write $X = X_1 \times X_2$ and similarly for Y , where X_i and Y_i are inner ideals in A^d . The set $Z_0(k)$ consists of pairs $X' \subset Y'$ where X' and Y' are inner ideals in A^d of dimension 1 and 10 respectively. Define $f: Z \rightarrow Z_0$ by $f(X \subset Y) = (X_1 \subset Y_1)$; it is G -equivariant.

On the other hand, a pair $X' \subset Y'$ in $Z_0(k)$ is the image of $(X', \psi(Y')) \subset (Y', \psi(X'))$ under f . Indeed, $\psi(Y')$ and $\psi(X')$ are inner ideals of the appropriate dimension by 3.33, and $X' \subset Y'$ implies $\psi(Y') \subset \psi(X')$.

The second and third lines of the table follow by similar reasoning.

Now let X be as in the last line of the table. The k -points of Z_0 are the 6-dimensional singular subspaces of A^d . Again writing X as (X_1, X_2) , we define f by $f(X) = X_1$. The equation $[X, X, V] = X$ is equivalent to the equations $\{X_i, X_{i+1}, A^d\} = X_i$ for $i = 1, 2$. Clearly, X_{i+1} is a subset of $\psi(X_i)$. But $\psi(X_i)$ also has dimension 6 by 3.33, hence $X_{i+1} = \psi(X_i)$. That is, X_2 is determined by X_1 . The conclusion now follows as in the case of the first line. \square

We now use the description of the projective homogeneous varieties of $G(A, K)$ given in the proposition to give a concrete criterion for determining whether $G(A, K)$ is isotropic.

5.3. Corollary. *For a group $G(A, K)$, the following are equivalent:*


- (a) $G(A, K)$ is isotropic.

- (b) *The projective homogeneous variety $\{\alpha_1, \alpha_6\}$ has a k -point.*
- (c) *There is a nonzero $x \in A_K$ such that*

$$x^\sharp = 0 \quad \text{and} \quad x \in \iota(x) \times A_K.$$

If these equivalent conditions hold, then the class $a(G(A, K))$ defined in 2.1 is a symbol in $H^3(k, 2)$.

Proof. We first prove that (a) implies (b); we suppose that $G(A, K)$ is isotropic. If K is $k \times k$, then we claim that the group is split or has the index displayed in 2.8. The only other index listed in [Ti 66b, p. 58] is



but this cannot be the index of a group with trivial Tits algebras [Ti 71, 6.4.5]. The claim is proved. Note that $a(G(A, K))$ is a symbol in $H^3(k, 2)$.

If K is a field, then $a(G(A, K))$ belongs to $H^3(k, 2)$ by Prop. 2.9. Therefore, $a(G(A, K))$ is a symbol by 4.8 and the index of $G(A, K)$ is in the top three lines of Table 2.4.

Summarizing the two preceding paragraphs, we find that the vertices α_1 and α_6 at the ends of the long arms of the Dynkin diagram are circled in the index, so the projective homogeneous variety $\{\alpha_1, \alpha_6\}$ has a k -point [BT, 6.4], i.e., (b) holds. The converse is standard: proper parabolic subgroups contain nontrivial split tori.

We now prove the equivalence of (b) and (c). By Prop. 5.2, (b) holds if and only if there are inner ideals $X \subset Y$ in A_K of ranks 1 and 10 such that $[X, Y, A_K] = 0$. Suppose first that such inner ideals $X \subset Y$ exist. We have $X = Kx$ for some nonzero $x \in A_K$ satisfying $x^\sharp = 0$ and

$$\{X, \iota(Y), A_K\} = [X, Y, A_K] = 0.$$

Since (3.30) implies (3.31), $\iota(Y)$ is contained in $\psi(X) = x \times A_K$. Comparing ranks using 3.33, we conclude that $X \subseteq Y = \iota(x) \times A_K$. In particular, (c) holds.

Conversely, suppose that (c) holds. Then $X := Kx$ and $Y := \iota(x) \times A_K$ are inner ideals of A_K of the desired ranks such that $X \subset Y$. Also, $[x, \iota(x) \times A_K, A_K] = \{x, x \times A_K, A_K\}$, which is zero by (3.10) and (3.11), so (b) holds. \square

5.4. Example. *If A has nonzero nilpotent elements, then $G(A, K)$ is isotropic. Indeed, A has a nonzero element x such that $x^\sharp = 0$ and $T(x) = 0$ by 3.26. Then $\iota(x) \times (-1) = x$ by (3.3).*

This example shows that in Theorem 0.2, (4) implies (1).

We now resolve an open question from [V 69].

5.5. Example. Consider again a group G as in Example 4.9, and write K for the associated quadratic extension. The group is of the form $G(\mathcal{T})$ for

a triple $\mathcal{T} = (V, K, P)$ where \mathcal{T} contains an “ α_2 -space”, i.e., an inner ideal X of V such that $\dim X = 6$ and $[X, X, V] = X$.

Over K , $\mathcal{T} \otimes K$ is isomorphic to a triple $\mathcal{T}(A, K \times K)$ for some Albert K -algebra A . However, A contains a 6-dimensional singular subspace, so it is split [Ra 77, Th. 1]. That is, $\mathcal{T} \otimes K$ is isomorphic to the split triple $\mathcal{T}(A_K^d, K \times K)$. (Alternatively, one can see this using Tits’s list of possible indexes for groups of type 1E_6 .) By Galois descent, we may identify V with

$$\{(a, tia) \mid a \in A_K^d\},$$

where t is some K -linear transformation of A_K^d . (Our ti is Veldkamp’s T .) As in the proof of Prop. 5.2, we have $tiX = \psi(X)$, where we have identified X with its first component in $A_K \times A_K$. For $x \in X$, we have $T(x, tix) = 0$ by [CG, 13.19].

Veldkamp was concerned with geometries whose points were 1-dimensional singular K -subspaces of A_K^d . In his language, the last sentence of the previous paragraph shows that Kx is a weakly isotropic point when x is nonzero. Veldkamp asked on page 291 of [V 69] if it is possible to have a weakly isotropic point and no strongly isotropic point. We now observe that there is no strongly isotropic point in this example. Suppose that $z \in A_K^d$ is nonzero and strongly isotropic, i.e., $\{z, tiz, A_K^d\}$ is zero. In that case, taking $Z = Ktiz$ and $Y = z \times A_K^d$ we find a pair of inner ideals of dimension 1 and 10 in \mathcal{T} such that $Z \subset Y$ and $[Z, Y, V] = 0$. But this contradicts the hypothesis on the index of $G(\mathcal{T})$, so no such z can exist, i.e., we have produced an explicit example of a situation where there is a weakly isotropic point and no strongly isotropic point.

6. EMBEDDINGS OF $k \times K$

In this section, we assume that K is a separable quadratic field extension of k , and write ι for its nontrivial k -automorphism. We fix a representative $\delta \in k^\times$ for the discriminant of K/k . (We take the naive definition of discriminant from, say, [Lang]. Consequently, in characteristic 2, we can and do take $\delta = 1$.) The purpose of this (long) section is to prove the following proposition, which in turn shows that (2) implies (3) in the main theorem, see 6.2.

6.1. Proposition. *Let A be an Albert k -algebra. Then $k \times K$ is a subalgebra of A if and only if A is isomorphic to $\mathcal{H}_3(C, \langle r, 1, \delta N_K(s) \rangle)$ for some octonion algebra C and elements $r \in k^\times$ and $s \in K^\times$ such that $T_K(s) \neq 0$.*

The assumption that K is a field is harmless. If K is $k \times k$, then we can take $\delta = 1$ and the proposition is easily seen to hold so long as k is not the field with two elements.

6.2. Proof that (2) implies (3). Before proceeding with the proof of the proposition, we first observe that it shows that (2) implies (3) in the main theorem. Prop. 6.1 gives that A is isomorphic to $\mathcal{H}_3(C, \langle r, 1, \delta N_K(s) \rangle)$.

That is, $f_3(A)$ is $[C]$ and $f_5(A)$ equals $[C] \cdot (-r) \cdot (-\delta N_K(s))$ by (3.25). The 2-Pfister bilinear form $\gamma := \langle\langle -r, -\delta N_K(s) \rangle\rangle$ satisfies (3). \square

6.3. Proof that (4) implies (2). The proposition also gives a proof that (4) implies (2) in the main theorem. Specifically, if A has a nilpotent, it is isomorphic to $\mathcal{H}_3(C, \langle -1, 1, \delta N_K(s) \rangle)$ for every $\delta \in k^\times$ and every $s \in K^\times$ with nonzero trace, since both algebras have the same invariants $f_3 = [C]$ and $f_5 = 0$. Then A contains $k \times K$ by the proposition.

6.4. Proof of 6.1: the easy direction. We suppose that A is $\mathcal{H}_3(C, \Gamma)$ for Γ as in the proposition and produce an explicit embedding of $k \times K$ in A . Fix $t \in K^\times$ such that $t^2 = \delta$, which implies

$$(6.5) \quad T_K(t) = 0 \quad \text{and} \quad N_K(t) = -\delta,$$

and define a map $\varphi: k \times K \rightarrow A$ by

$$\begin{aligned} \varphi(\alpha, a) := & \alpha e_{11} + \\ & T_K(s)^{-1} \left(N_K(s, a) e_{22} + N_K(s, \iota(a)) e_{33} + t^{-1}(\iota(a) - a) 1_C [23] \right) \end{aligned}$$

for $\alpha \in k, a \in K$. Note that $t^{-1}(\iota(a) - a)$ is in k in all characteristics by (6.5), i.e., the image of φ really is in A and not just in A_K . Also, φ sends 1 to 1. Hence it suffices to show that φ preserves norms. Plugging in to the explicit formula for the norm on J from (3.16), we find

$$N[\varphi(\alpha, a)] = \frac{\alpha}{T_K(s)^2} \left(N_K(s, a) N_K(s, \iota(a)) - N_K(s) (\iota(a) - a)^2 \right).$$

Writing out, for example, $N_K(s, a)$ as $s\iota(a) + \iota(s)a$, we find that $N[\varphi(\alpha, a)]$ is $\alpha N_K(a)$ as desired. \square

The converse implication of Prop. 6.1 takes a bit more work. Fortunately, we can rely on the proof of [PR84, Th. 3.2] to simplify the task. For the sake of clarity, we still indicate the main steps of that proof insofar they are relevant for our purposes.

6.6. Proof of 6.1: the difficult direction. Assume now that $E := k \times K$ is a subalgebra of A as in the statement of Prop. 6.1. We let ι act on $A_K := A \otimes K$ and $E_K := E \otimes K \subset A_K$ through the second factor. Since A inherits zero divisors from E , it is reduced. We write C for its coordinate algebra, forcing $C_K := C \otimes K$ to be the coordinate algebra of A_K .

Step 1. A coordinatization for E_K . Since E_K is split but E is not (because K is a field by hypothesis), there is a frame (e_1, e_2, e_3) of A_K satisfying

$$E_K = Ke_1 \oplus Ke_2 \oplus Ke_3 \quad \text{with} \quad \iota(e_1) = e_1, \quad \iota(e_2) = e_3, \quad \iota(e_3) = e_2.$$

This setup will remain fixed till the end of the proof. It implies

$$E = \{ \alpha e_1 + a e_2 + \iota(a) e_3 \mid \alpha \in k, a \in K \}.$$

Step 2. A coordinatization for A_K . Following [PR 84, Lemma 3.5], we find a coordinatization of A_K having the form

$$(6.7) \quad A_K = \mathcal{H}_3(C_K, \langle 1, s, \iota(s) \rangle) \quad \text{for some } s \in K^\times$$

such that $e_i = e_{ii}$ for all i , the trace $T_K(s)$ is not zero, and there is an ι -semilinear involution τ_K of C_K that commutes with the conjugation of C_K and satisfies the relation

$$\iota(x) = \iota(a_1)e_1 + \iota(a_3)e_2 + \iota(a_2)e_3 + \tau_K(x_1)[23] + \tau_K(x_3)[31] + \tau_K(x_2)[12]$$

for all $x = \sum a_i e_{ii} + \sum x_i [jl] \in A_K$. In particular,

$$(6.8) \quad A = \left\{ \begin{array}{l} \alpha e_1 + a e_2 + \iota(a) e_3 \\ + x_1 [23] + x_2 [31] + \tau_K(x_2) [12] \end{array} \middle| \begin{array}{l} \alpha \in k, a \in K, \\ x_1 \in C_K^{\tau_K}, x_2 \in C_K \end{array} \right\},$$

where $C_K^{\tau_K}$ denotes the subspace of C_K consisting of elements fixed by τ_K .

Step 3. A frame for A . By [PR 84, Lemma 3.6], $F = (d_1, d_2, d_3)$, where

$$(6.9) \quad \begin{aligned} d_1 &:= e_{11}, \\ d_2 &:= T_K(s)^{-1}(s e_{22} + \iota(s) e_{33} + 1[23]) \\ d_3 &:= T_K(s)^{-1}(\iota(s) e_{22} + s e_{33} - 1[23]), \end{aligned}$$

is a frame of A . Furthermore, the map

$$(6.10) \quad \tau: C \rightarrow C \text{ defined by } x \mapsto \tau(x) := \overline{\tau_K(x)},$$

is an ι -semilinear automorphism of C_K , forcing $B := C_K^\tau$ to be a composition algebra over k such that $C_K = B \otimes K$.

Step 4. Peirce components of A . As in 6.4, we now fix $t \in K^\times$ satisfying $t^2 = \delta$, hence (6.5). Then the following relations hold:

$$(6.11) \quad A_{12}(F) = \{x_2[31] + \overline{x_2}[12] \mid x_2 \in B\},$$

$$(6.12) \quad A_{31}(F) = \{(st)x_2[31] + \iota(st)\overline{x_2}[12] \mid x_2 \in B\}.$$

While (6.11) was established in [PR 84, Lemma 3.6b], the proof of (6.12) is a bit more involved and runs as follows. Standard facts about Peirce components and (6.8), (6.9) imply

$$\begin{aligned} A_{31}(F) &= A_1(d_3) \cap A_1(d_1) = A_1(d_3) \cap A \cap (C_K[31] + C_K[12]) \\ &= A_1(d_3) \cap \{x_2[31] + \tau_K(x_2)[12] \mid x_2 \in C_K\}. \end{aligned}$$

For $x_2 \in C_K$, the element $x := x_2[31] + \tau_K(x_2)[12]$ satisfies

$$\begin{aligned} T_K(s)(d_3 \times x) &= (\iota(s)e_{22} + s e_{33} - 1[23]) \times (x_2[31] + \tau_K(x_2)[12]) \\ &= -(\overline{s\tau_K(x_2)} + \iota(s)x_2)[31] - (\iota(s)\overline{x_2} + s\tau_K(x_2))[12] \end{aligned}$$

by (3.20) and (6.7), and x belongs to $A_1(d_3)$ if and only if this expression is zero (Lemma 3.14a). Since $\iota(t) = -t$ by (6.5), we may apply (6.10) to conclude that this in turn is equivalent to the equation $st\tau(x_2) = \iota(st)x_2$, i.e., to $x_2 \in (st)B$.

Step 5. Conclusion of proof. Using (3.21) and (6.7), it is straightforward to check the relations

$$(6.13) \quad S(x_2[31] + \overline{x_2}[12]) = -T_K(s)N_B(x_2),$$

$$(6.14) \quad S((st)x_2[31] + \iota(st)\overline{x_2}[12]) = -T_K(s)N_K(s)\delta N_B(x_2)$$

for all $x_2 \in B$. Following [McC 66, p. 1077], we now choose a coordinatization of A that has the form

$$A = \mathcal{H}_3(C, \Gamma^0) \text{ for } \Gamma^0 = \langle \gamma_1^0, 1, \gamma_3^0 \rangle \in \text{GL}_3(k),$$

and that matches F with the diagonal frame of $\mathcal{H}_3(C, \Gamma^0)$. Comparing (6.13), (6.14) with (3.22), we conclude

$$(6.15) \quad \langle -\gamma_1^0 \rangle N_C \cong S|_{A_{12}(F)} \cong \langle -T_K(s) \rangle N_B,$$

$$(6.16) \quad \langle -\gamma_3^0 \gamma_1^0 \rangle N_C \cong S|_{A_{31}(F)} \cong \langle -T_K(s)N_K(s)\delta \rangle N_B.$$

In particular, the composition algebras B, C over k have similar norm forms and hence are isomorphic, allowing us to identify $B = C$ from now on. Then (6.15) and (6.16) imply that A and $\mathcal{H}_3(C, \langle T_K(s), 1, \delta N_K(s) \rangle)$ have the same mod-2 invariants, hence are isomorphic. \square

7. THE CASE WHERE $T(x) = 0$

In this and the following section, we will prove that (1) implies (2) in the main theorem. Suppose that (1) holds, so that by Cor. 5.3 A_K contains a nonzero element x satisfying

$$(7.1) \quad x^\sharp = 0 \quad \text{and} \quad x \in \iota(x) \times A_K.$$

In this section, we treat the case where x satisfies (7.1) and has trace zero. The next section treats the case where the trace of x is nonzero.

7.2. Proposition. *Let A be an Albert k -algebra. There is a nonzero element $x \in A_K$ satisfying*

$$x^\sharp = 0, \quad x \in \iota(x) \times A_K, \quad \text{and} \quad T(x) = 0$$

if and only if A contains nonzero nilpotent elements.

Proof. We suppose that A_K contains an x as in the statement of the proposition; the other direction was treated in Example 5.4. If K is equal to $k \times k$, then $A_K = A \times A$ and $x = (x_1, x_2)$ for $x_i \in A$ not both zero with $x_i^\sharp = 0$ and $T(x_i) = 0$. Therefore one of the elements $x_1, x_2 \in A$ is a nonzero nilpotent by 3.26. We are left with the case where K is a field.

For sake of contradiction, we further assume that A has no nonzero nilpotents. Write $K = k[d]$ where $d \in K$ has trace 1. Set $\delta := N_K(d) \in k^\times$, so that $d^2 = d - \delta$. Write

$$x = y + dz \quad \text{with } y, z \in A \text{ not both zero.}$$

Because $x^\sharp = 0$, we have:

$$(7.3) \quad y^\sharp = \delta z^\sharp \quad \text{and} \quad z^\sharp = -y \times z.$$

Clearly,

$$(7.4) \quad T(y) = T(z) = 0.$$

We first argue that neither y nor z is invertible in A . By (3.2) we have:

$$(7.5) \quad N(y)y = (y^\sharp)^\sharp = \delta^2(z^\sharp)^\sharp = \delta^2 N(z)z.$$

On the other hand, comparing

$$\begin{aligned} y^\sharp \times (y \times z) &= N(y)z + T(y^\sharp, z)y && \text{(by (3.7))} \\ &= N(y)z + \delta T(z^\sharp, z)y && \text{(by (7.3))} \\ &= N(y)z + 3\delta N(z)y && \text{(by (3.8))} \end{aligned}$$

with

$$\begin{aligned} y^\sharp \times (y \times z) &= -\delta z^\sharp \times z^\sharp && \text{(by (7.3))} \\ &= -2\delta z^{\sharp\sharp} = -2\delta N(z)z, && \text{(by (3.2))} \end{aligned}$$

we obtain

$$(7.6) \quad -2\delta N(z)z = N(y)z + 3\delta N(z)y.$$

Similarly,

$$\begin{aligned} \delta z^\sharp \times (z \times y) &= \delta N(z)y + \delta T(z^\sharp, y)z && \text{(by (3.7))} \\ &= \delta N(z)y + T(y^\sharp, y)z && \text{(by (7.3))} \\ &= \delta N(z)y + 3N(y)z && \text{(by (3.8))} \end{aligned}$$

and

$$\begin{aligned} \delta z^\sharp \times (z \times y) &= -\delta z^\sharp \times z^\sharp && \text{(by (7.3))} \\ &= -2\delta z^{\sharp\sharp} = -2\delta N(z)z && \text{(by (3.2))} \end{aligned}$$

imply

$$(7.7) \quad -2\delta N(z)z = 3N(y)z + \delta N(z)y.$$

Subtracting (7.6) from (7.7), we conclude $2N(y)z = 2\delta N(z)y$, which implies

$$(7.8) \quad N(y)z = \delta N(z)y$$

for char $k \neq 2$, while this follows directly from (7.6) for char $k = 2$. Thus (7.8) holds in full generality. By (7.5), assuming that one of the elements y, z is invertible, both are, and

$$\begin{aligned} N(y)^2 z^\sharp &= (N(y)z)^\sharp = (\delta N(z)y)^\sharp && \text{(by (7.8))} \\ &= \delta^2 N(z)^2 y^\sharp = \delta^3 N(z)^2 z^\sharp && \text{(by (7.3))} \end{aligned}$$

implies

$$(7.9) \quad N(y)^2 = \delta^3 N(z)^2.$$

On the other hand,

$$\begin{aligned}\delta N(y)z^\sharp &= -\delta y \times [N(y)z] && \text{(by (7.3))} \\ &= -\delta^2 N(z)y \times y && \text{(by (7.8))} \\ &= -2\delta^2 N(z)y^\sharp = -2\delta^3 N(z)z^\sharp, && \text{(by (7.3))}\end{aligned}$$

which yields

$$(7.10) \quad N(y) = -2\delta^2 N(z).$$

Hence $\text{char } k \neq 2$, and comparing the square of (7.10) with (7.9) shows $\delta = \frac{1}{4}$. But then the minimum polynomial of d over k becomes $X^2 - X + \frac{1}{4} = (X - \frac{1}{2})^2$, contradicting the fact that K is a field. We have thus established the relation

$$(7.11) \quad N(y) = N(z) = 0.$$

By assumption, A does not contain nonzero nilpotent elements. Hence (7.4), (7.11) imply that $S(y), S(z)$ cannot both be zero since, otherwise, y, z were both nilpotent by (3.27). But since $S(y) = \delta S(z)$ by (7.3), we conclude $S(y)$ and $S(z)$ are both nonzero. On the other hand, (7.11) combines with (3.2) to yield $y^{\sharp\sharp} = z^{\sharp\sharp} = 0$, forcing

$$e = S(y)^{-1}y^\sharp = S(z)^{-1}z^\sharp \quad \text{(by (7.3))}$$

to be a primitive idempotent of A . The relation

$$e \times x = S(y)^{-1}(y \times y^\sharp) + dS(z)^{-1}(z \times z^\sharp)$$

combined with (3.9), (7.4), and (7.11) gives:

$$e \times x = -(y + dz) = T(x)(1 - e) - x.$$

Lemma 3.14a shows that x belongs to $(A_K)_0(e)$, hence so does $\iota(x)$.

Now fix $v \in A_K$ such that $x = \iota(x) \times v$ and write $v = ae + v_1 + v_0$ for $a \in K$ and $v_i \in (A_K)_i(e)$ with $i = 0, 1$. Since $\iota(x)$ is in $(A_K)_0(e)$, we may apply Lemma 3.14a again to conclude $\iota(x) \times (ae) = -a\iota(x) \in (A_K)_0(e)$. On the other hand, Lemma 3.14c gives $\iota(x) \times v_0 = S(\iota(x), v_0)e \in (A_K)_2(e)$. Finally, using the circle product $a \circ b := \{a, 1, b\} = U_{a,b}1$, we obtain

$$\iota(x) \times v_1 = \iota(x) \circ v_1 - T(\iota(x))v_1 - T(v_1)\iota(x) + [T(\iota(x))T(v_1) - T(\iota(x), v_1)]1$$

by (3.5) and (3.6) linearized. But, as the Peirce decomposition is orthogonal relative to the generic trace, this is just $\iota(x) \circ v_1$, which is in $(A_K)_1(e)$. Decomposing $a = \alpha + \beta d$ with $\alpha, \beta \in k$ and comparing Peirce components of $x = \iota(x) \times v$ relative to e , a short computation yields

$$y + dz = \iota(x) \times v = -a\iota(x) = -(\alpha y + (\alpha + \delta\beta)z) - d(\beta y - \alpha z),$$

hence

$$(7.12) \quad (1 + \alpha)y = -(\alpha + \delta\beta)z \quad \text{and} \quad (1 - \alpha)z = -\beta y.$$

To complete the proof, it suffices to show $z = -2y$, because this implies $0 = x^\sharp = (1 - 2d)^2 y^\sharp$, hence $2d = 1$, a contradiction. Now, if $\beta = 0$, then $\alpha = 1$ and $2y = -z$ by (7.12). But if $\beta \neq 0$, then

$$\begin{aligned} \beta z^\sharp &= -\beta y \times z = (1 - \alpha)z \times z && \text{(by (7.3) and (7.12))} \\ &= 2(1 - \alpha)z^\sharp, \end{aligned}$$

which yields $\beta = 2(1 - \alpha)$, hence $\beta z = 2(1 - \alpha)z = -2\beta y$ by (7.12), and again we end up with $z = -2y$. \square

8. THE CASE WHERE $T(x) \neq 0$

We now treat the other possible consequence of hypothesis (1).

8.1. Proposition. *There is a nonzero element $x \in A_K$ satisfying*

$$x^\sharp = 0, \quad x \in \iota(x) \times A_K, \quad \text{and} \quad T(x) \neq 0$$

if and only if $k \times K$ is a subalgebra of A .

Proof. First suppose that K is not a field, i.e., $K = k \times k$. Then A_K is identified with $A \times A$ and ι acts via the switch. If A_K contains an element $x = (x_1, x_2)$ as in the statement of the proposition, then one of the elements $x_1, x_2 \in A$ is singular, hence A is reduced. It follows that $k \times k \times k$ is a subalgebra. Conversely, if $k \times k \times k$ is a subalgebra of A , then the standard basis vectors e_1, e_2, e_3 in k^3 form a complete orthogonal system of primitive idempotents in A . Putting $x = (e_1, e_2)$, we find that $x^\sharp = 0$, x has trace one, and

$$\iota(x) \times e_3 = (e_2 \times e_3, e_1 \times e_3) = x.$$

We are left with the case where K is a field. Suppose first that A_K contains an element x as in the statement of the proposition. Then putting $b := T(x) \in K^\times$, the element $e := b^{-1}x$ is a primitive idempotent in A_K and

$$\iota(e) = \iota(b)^{-1}\iota(x) \in \iota(b)^{-1}(x \times A_K) = e \times (\iota(b)^{-1}b A_K) \subseteq e \times A_K.$$

But since $e \times e = 2e^\sharp = 0$ and $f = 1 - e$ is in $(A_K)_0(e)$, Lemma 3.14a implies that the map $x \mapsto e \times x$ kills $(A_K)_2(e) + (A_K)_1(e)$ and stabilizes $(A_K)_0(e)$. So $\iota(e)$ is in $(A_K)_2(e)$ and $e, \iota(e), c$ are orthogonal primitive idempotents in A_K where $c := 1 - e - \iota(e)$. The idempotent c , remaining fixed under ι , belongs to A . Thus

$$k \times K \rightarrow A \quad \text{via} \quad (\alpha, a) \mapsto \alpha c + ae + \iota(a)\iota(e),$$

is an embedding of Jordan algebras over k .

Conversely, suppose that K is a field and $k \times K$ is a subalgebra of A . By hypothesis, $A \otimes K$ contains $K \times (K \otimes K) = K \times K \times K$ as a subalgebra whose unit vectors u_1, u_2, u_3 form a complete orthogonal system of primitive idempotents in $A \otimes K$ such that u_1 is in A and ι interchanges u_2 and u_3 . Thus $u_2 \in A \otimes K$ is a singular element with trace 1 that satisfies

$$u_2 = u_3 \times u_1 \in \iota(u_2) \times A_K. \quad \square$$

9. SUFFICIENT CONDITION FOR ISOMORPHISM

Let A be a reduced Albert algebra over k with coordinate algebra C . By 3.24, the 5-Pfister form corresponding to $f_5(A)$ may be written as $N_C \otimes \gamma$ for some 2-Pfister bilinear form γ . Since reduced Albert algebras are classified by their mod-2 invariants, we can write A as $\mathcal{H}_3(C, \gamma)$. By the same token, given octonion algebras C, C' and 2-Pfister bilinear forms γ, γ' over k , the algebras $\mathcal{H}_3(C, \gamma)$ and $\mathcal{H}_3(C', \gamma')$ are isomorphic if and only if $N_C \cong N_{C'}$ and $N_C \otimes \gamma \cong N_{C'} \otimes \gamma'$. The goal of this section is to prove a weak analogue of this statement for hermitian Albert triples.

9.1. Proposition. *Let K be a quadratic étale k -algebra, let C be an octonion k -algebra, and let γ and γ' be 2-Pfister bilinear forms. If $\gamma \otimes N_K$ is isomorphic to $\gamma' \otimes N_K$, then $\mathcal{T}(\mathcal{H}_3(C, \gamma), K)$ is isomorphic to $\mathcal{T}(\mathcal{H}_3(C, \gamma'), K)$.*

It is natural to wonder if a stronger result holds, namely if $N_C \otimes N_K \otimes \gamma$ is isomorphic to $N_C \otimes N_K \otimes \gamma'$, are the triples $\mathcal{T}(\mathcal{H}_3(C, \gamma), K)$ and $\mathcal{T}(\mathcal{H}_3(C, \gamma'), K)$ necessarily isomorphic? (The tensor products in the question only make sense in characteristic different from 2.) The answer is no, as Example 12.2 below shows.

The proposition will follow easily (see the end of this section) from an alternative construction of hermitian Albert triples that we now describe. First we claim that the general linear group $\mathrm{GL}_3(k)$ acts on the split Albert algebra $A^d = \mathcal{H}_3(C^d, 1)$ via

$$\phi_g(j) := gjg^t \quad (g \in \mathrm{GL}_3(k), j \in A^d)$$

in such a way that

$$(9.2) \quad N(\phi_g(j)) = (\det g)^2 N(j).$$

To see this, it suffices to show $gA^d g^t \subseteq A^d$ and (9.2) for *elementary* matrices $g \in \mathrm{GL}_3(k)$, which follows easily by brute force using (3.16). (This is the argument given in [J61, §5].) Also, we have $\phi_g^\dagger = \phi_{g^{-t}}$ by the argument in [J61, p. 77].

In particular, the map ϕ_g , for $g \in \mathrm{SL}_3(k)$, is an automorphism of the split hermitian Albert triple \mathcal{T}^d via

$$\phi_g \cdot (j_1, j_2) = (gj_1 g^t, g^{-t} j_2 g^{-1}).$$

This gives a map

$$(9.3) \quad G_2 \times (\mathrm{SL}_3 \times \mathbb{Z}/2\mathbb{Z}) \rightarrow \mathrm{Aut}(\mathcal{T}^d),$$

and a corresponding map

$$(9.4) \quad H^1(k, G_2) \times H^1(k, \mathrm{SL}_3 \times \mathbb{Z}/2\mathbb{Z}) \rightarrow H^1(k, \mathrm{Aut}(\mathcal{T}^d)).$$

This last map takes an octonion k -algebra C , a quadratic étale k -algebra K , and a rank 3 K/k -hermitian form as inputs, and it gives a hermitian Jordan triple as output.

9.5. Lemma. *For every quadratic étale k -algebra K , octonion k -algebra C , and 3-dimensional symmetric bilinear form γ , construction (9.4) sends the hermitian form deduced from γ to $\mathcal{F}(\mathcal{H}_3(C, \gamma), K)$.*

Sketch of proof. The proof is essentially the same as the proof of [Fe 78, Prop. 3.2], so we omit it. One observes that construction (9.4) sends the hermitian form to $\mathcal{F}(A, K)$, where A is the γ -isotope of $\mathcal{H}_3(C, 1)$. But A is isomorphic to $\mathcal{H}_3(C, \gamma)$ by [J 68, p. 61, Th. 14] or [McC 66, p. 1077]. \square

9.6. Example. In [V 68] and [V 69], Veldkamp considers groups of type 2E_6 constructed by (9.4) from an octonion k -algebra C and a cocycle in $Z^1(K/k, \mathrm{SL}_3 \rtimes \mathbb{Z}/2\mathbb{Z})$ whose value at ι is $(\langle 1, 1, -1 \rangle, 1)$. This cocycle corresponds to the K/k -hermitian form deduced from $\langle 1, 1, -1 \rangle$. By the lemma, such a group is isomorphic to $G(\mathcal{H}_3(C, \langle 1, 1, -1 \rangle), K)$, i.e., is one of the groups in the first three lines of Table 2.4.

Proof of Proposition 9.1. Under our hypotheses, the rank 3 K/k -hermitian forms deduced from the pure parts of γ and γ' are isometric. Therefore, the hermitian Jordan triples constructed from them via (9.4) are isomorphic. Now apply Lemma 9.5. \square

9.7. Proof that (3) implies (1). Suppose that (3) holds in the statement of Theorem 0.2. Then A is isomorphic to $\mathcal{H}_3(C, \gamma)$ where $\gamma \otimes N_K$ is hyperbolic. By Prop. 9.1, $G(A, K)$ is isomorphic to $G(\mathcal{H}_3(C, \langle 1, -1, 1 \rangle), K)$, and (1) holds by Example 5.4. \square

9.8. Vista. The center of SL_3 is identified with the center of the split group $E_6 := \mathrm{Aut}(\mathcal{T}^d)$ via (9.3), and this gives a map

$$G_2 \times \mathrm{Aut}(\mathrm{PSL}_3) \rightarrow \mathrm{Aut}(E_6).$$

This leads to a construction of groups of type E_6 (with possibly non-trivial Tits algebras), with inputs an octonion algebra, say C , and a central simple associative algebra of degree 3 with unitary involution fixing k , say (B, τ) . Tits's other construction of Lie algebras of type E_6 (corresponding to the E_6 in the bottom row of the “magic square” [J 71, p. 98]) uses identical inputs, and it is natural to guess that it produces the Lie algebra of the group arising from the construction above. In any case, one can ask: *What is the index of the group constructed from given C and (B, τ) ?*

We remark that some of the techniques in this paper can be adapted to attack this question. For §4, hermitian Jordan triples should be replaced by a new type of algebraic structure in a manner analogous to the replacement of quadratic forms by algebras with orthogonal involution as in [KMRT, §5]. The description of the homogenous projective varieties in §5 can be translated to this new structure. We leave the details to the interested reader.

10. THE CASE WHERE A IS SPLIT BY K

10.1. In the notation of the main theorem, we always have (4) implies (1) by Example 5.4. Conversely, suppose that K splits A and (3) holds. In particular, A is reduced and we may write $f_3(A)$ as $[K] \cdot \tau$ for some 2-Pfister bilinear form τ , hence

$$f_5(A) = f_3(A) \cdot \gamma = \tau \cdot 0 = 0.$$

That is, A has nonzero nilpotents. This proves the final sentence of Theorem 0.2. \square

We now provide an example to show that one really needs the hypothesis that K splits A in the implication that (3) implies (4).

10.2. Example. Take $k = \mathbb{Q}(x, y, z, u, d)$ to be the rational function field in five variables over \mathbb{Q} , let C be the octonion k -algebra corresponding to the 3-Pfister form $\langle\langle x, y, z \rangle\rangle$, and put

$$A := \mathcal{H}_3(C, \langle d, 1, -u \rangle) \quad \text{and} \quad K := k(\sqrt{d}).$$


Then $f_3(A) = (x) \cdot (y) \cdot (z)$ and $f_5(A) = f_3(A) \cdot \gamma$ for $\gamma = \langle\langle -d, u \rangle\rangle$. Since $(-d) \cdot (d) = 0$, we have $\gamma \cdot (d) = 0$, i.e., (3) holds. On the other hand, the 5-Pfister $\langle\langle x, y, z, -d, u \rangle\rangle$ is anisotropic by Springer's Theorem [Lam, §VI.1], so $f_5(A)$ is nonzero and A contains no nonzero nilpotents.

11. APPLICATIONS TO ISOTROPIC GROUPS

In Example 4.6, we gave some examples of fields over which every group of type E_6 with trivial Tits algebras is—up to isogeny—of the form $G(A, K)$. For *isotropic* groups, there is a statement that holds for every field.

11.1. Proposition. *The map $(C, K) \mapsto G(\mathcal{H}_3(C, \langle 1, 1, -1 \rangle), K)$ defines a bijection between isomorphism classes of*

- (a) *pairs (C, K) where C is an octonion k -algebra and K is a quadratic étale k -algebra; and*
- (b) *simply connected groups of type E_6 over k that are quasi-split or whose index is one of*



Furthermore, the groups in (b) are precisely the isotropic groups of the form $G(A, K)$ for some Albert k -algebra A and some quadratic étale k -algebra K .

The proof amounts to collecting results from earlier sections. Note that $\mathcal{H}_3(C, \langle 1, 1, -1 \rangle)$ has nonzero nilpotents, so the map indeed produces groups as in (b) by Cor. 5.3 and Example 5.4.

Proof. Write f for the map described in the proposition. Write g for the map that takes a group G as in (b) and gives the pair (C, K) where C is the octonion k -algebra corresponding to $a(G)$ (a symbol in $H^3(k, 2)$ by

Cor. 5.3) and K/k is $k \times k$ in case G has type 1E_6 and is the separable quadratic extension of k over which the Galois group acts trivially on the Dynkin diagram of G in case G has type 2E_6 . The composition gf is the identity by 4.8, so f is injective and g is surjective. On the other hand, g is injective by 2.8 and Prop. 2.9. Since gfg equals g and g is an isomorphism, we conclude that gf is the identity. This proves that f is an isomorphism.

We now prove the “furthermore”. We observed in the proof of Cor. 5.3 that, if $G(A, K)$ is isotropic, then it is one of the groups in (b). Conversely, every group in (b) is of the form $G(A, K)$ by the surjectivity of the map f . \square

11.2. A weaker version of Prop. 11.1 is: *Every group whose index appears in the first three rows of Table 2.4 is isomorphic to $G(\mathcal{H}_3(C, \langle 1, 1, -1 \rangle), K)$ for some octonion algebra C and some K .* This is approximately the content of [V 68, 3.3] and [V 69, 3.2].

As a corollary to the proposition, we obtain: *Conditions (1)–(3) in Th. 0.2 are equivalent to*


$$(11.3) \quad G(A, K) \text{ is isomorphic to } G(A', K) \text{ for some Albert } k\text{-algebra } A' \text{ with nonzero nilpotents.}$$

11.4. Example. Consider the groups G and G' over \mathbb{R} defined by

$$G := G(\mathcal{H}_3(\mathbb{O}, \langle 1, 1, -1 \rangle), \mathbb{C}) \quad \text{and} \quad G' := G(\mathcal{H}_3(\mathbb{O}, 1), \mathbb{C}),$$

where \mathbb{O} denotes the division octonion algebra over \mathbb{R} . The invariants $a(G)$ and $a(G')$ both equal the unique nonzero element in $H^3(k, 2)$ by 4.8, but G is isotropic and G' is anisotropic by the main theorem. (Said differently, the Killing form of G has signature -14 [J 71, p. 120] and G' is the compact simply connected E_6 .)

Note that the index of G is



by Prop. 2.3, but the Rost invariant of G has nonzero kernel. Specifically, for $\alpha \in H^1(k, G)$ such that G' is isomorphic to G twisted by α , $r_G(\alpha)$ is zero by 2.5, but α is not the zero class in $H^1(k, G)$ because G' is not isomorphic to G .

This example is in contrast with the quasi-split groups of type 2E_6 , for which the Rost invariant always has zero kernel by [Ga].

11.5. Example. We abuse notation by writing \mathbb{O} also for the unique octonion division algebra over \mathbb{Q} . Consider the groups G and G' over \mathbb{Q} defined by

$$G := G(\mathcal{H}_3(\mathbb{O}, \langle 1, 1, -1 \rangle), \mathbb{Q}(\sqrt{p})) \quad \text{and} \quad G' := G(\mathcal{H}_3(\mathbb{O}, 1), \mathbb{Q}(\sqrt{p})),$$

where p is a prime number. Again G' is anisotropic, and this time the group G has index



by the main theorem and Prop. 2.9, because $\mathbb{Q}(\sqrt{p})$ does not split \mathbb{O} . As in the preceding example, the Rost invariant r_G has nonzero kernel.

12. FINAL OBSERVATIONS

We close this paper by applying our main theorem to give an easy criterion for when $G(A, K)$ is isotropic over special fields.

12.1. Proposition. *Let k be a SAP field of characteristic zero such that $cd_2 k(\sqrt{-1}) \leq 2$. A group $G(A, K)$ is isotropic if and only if A is reduced and $f_5(A) \cdot [K] = 0$.*

A precise definition and basic properties of SAP (“Strong Approximation Property”) fields can be found in [EL]. Every algebraic extension of \mathbb{Q} (not necessarily of finite degree) and $\mathbb{R}((x))$ satisfy the hypothesis of the proposition. See [BP, pp. 653–655] for a summary of basic properties of fields k as in the proposition.

The restriction on the characteristic is harmless. The prime characteristic analogue of the proposition is a corollary of the statement: *If $H^5(k, 2)$ is zero and A is reduced, then $G(A, K)$ is isotropic.* To see this, note that the hypotheses imply that the subgroup $\text{Aut}(A)$ of $G(A, K)$ is itself isotropic.

Proof of the proposition. If $G(A, K)$ is isotropic, then A is reduced and $f_5(A) \cdot [K]$ is zero by the main theorem, so we assume that A is reduced and $f_5(A) \cdot [K]$ is zero and prove the converse. Because k is SAP, there is some $\gamma_1 \in k^\times$ that is positive at every ordering where $f_5(A)$ is zero and negative at every ordering where $f_5(A)$ is nonzero. Put $\gamma := \langle\langle -1, \gamma_1 \rangle\rangle$. Because $f_3(A)$ divides $f_5(A)$, it follows that $f_5(A)$ equals $f_3(A) \cdot \gamma$ and $\gamma \cdot [K]$ equals zero over every real-closure of k . The hypothesis on the cohomological dimension implies that those same equalities hold over k . We conclude that $G(A, K)$ is k -isotropic by the main theorem. \square

We observed in Prop. 9.1 that one can change γ somewhat without changing the isomorphism class of $G(\mathcal{H}_3(C, \gamma), K)$. Motivated by Prop. 12.1, one might hope that the main theorem still holds with (3) replaced by

$$(3') \quad A \text{ is reduced and } f_5(A) \cdot [K] = 0.$$

(We remark that the expression $f_5(A) \cdot [K]$ only makes sense when $\text{char } k \neq 2$.) Clearly, (3) implies (3'). We now give an explicit example where (3') holds but (3) does not. That is, such a replacement is not possible. We thank Detlev Hoffmann for showing us this example.

12.2. Example. Fix k_0 to be the purely transcendental extension $\mathbb{Q}(x, y, z, u, v, d)$ of the rationals (say). Let k denote the function field over k_0 of the 6-Pfister form $\langle\langle x, y, z, u, v, d \rangle\rangle$. Define a k -group $G(A, K)$ via

$$A := \mathcal{H}_3(C, \langle -u, -v, uv \rangle) \quad \text{and} \quad K := k(\sqrt{d}),$$

where C is the octonion k -algebra with norm $\langle\langle x, y, z \rangle\rangle$. Clearly, (3') holds over k .

On the other hand, take E to be the function field of $q := \langle\langle x, y, z \rangle\rangle' + \langle d \rangle$ over k , where $\langle\langle x, y, z \rangle\rangle'$ denotes the pure part of the 3-Pfister. Clearly, $f_3(A) = (x) \cdot (y) \cdot (z)$ is killed by EK , that is, $[K]$ divides $f_3(A)$ over E . We now argue that $f_5(A)$ is not zero over E , which will show that (3) fails over E hence also over k . Note that $f_5(A) = (x) \cdot (y) \cdot (z) \cdot (u) \cdot (v)$ is nonzero over k_0 and remains nonzero over k by Hoffmann's separation theorem [Lam, X.4.34].

For sake of contradiction, suppose that $f_5(A)$ is killed by E . Then by Cassels-Pfister [Lam, X.4.8], for $a \in k^\times$ represented by q and $b \in k^\times$ represented by $f_5(A)$, we have that $\langle ab \rangle q$ is a subform of $f_5(A)$. Taking $a = b = -x$, we find that q is a subform of $f_5(A)$ over k . Computing in the Witt ring of k , it follows that

$$\langle 1, -d \rangle + \langle\langle x, y, z \rangle\rangle \langle -u, -v, uv \rangle$$

is k -isotropic. But this form is k_0 -anisotropic and remains anisotropic over k because its dimension is $26 < 32 < 64 = \dim \langle\langle x, y, z, u, v, d \rangle\rangle$, again using the separation theorem. This is a contradiction, which shows that (3) does not hold.

Acknowledgements. The authors are indebted to Detlev Hoffmann, Ottmar Loos, Erhard Neher, and Michel Racine for useful discussions on the subject of this paper.

REFERENCES

- [AB] R. Aravire and R. Baeza, *Milnor's K-theory and quadratic forms over fields of characteristic 2*, Comm. Alg. **20** (1992), 1087–1107.
- [BP] E. Bayer-Fluckiger and R. Parimala, *Classical groups and the Hasse principle*, Ann. of Math. (2) **147** (1998), 651–693.
- [BT] A. Borel and J. Tits, *Groupes réductifs*, Inst. Hautes Études Sci. Publ. Math. **27** (1965), 55–150.
- [CG] M. Carr and S. Garibaldi, *Geometries, the principle of duality, and algebraic groups*, Expo. Math. **24** (2006), 195–234.
- [DG] M. Demazure and P. Gabriel, *Groupes algébriques. Tome I*, Masson, Paris, 1970.
- [Dy] E.B. Dynkin, *Semisimple subalgebras of semisimple Lie algebras*, Amer. Math. Soc. Transl. (2) **6** (1957), 111–244, [Russian original: Mat. Sbornik N.S. **30(72)** (1952), 349–462].
- [EL] R. Elman and T.Y. Lam, *Quadratic forms over formally real fields and pythagorean fields*, Amer. J. Math. **94** (1972), 1155–1194.
- [Fa] J.R. Faulkner, *Octonion planes defined by quadratic Jordan algebras*, Mem. Amer. Math. Soc. (1970), no. 104.
- [Fe 69] J.C. Ferrar, *Lie algebras of type E_6* , J. Algebra **13** (1969), 57–72.

- [Fe 78] ———, *Lie algebras of type E_6 . II*, J. Algebra **52** (1978), no. 1, 201–209.
- [Ga] R.S. Garibaldi, *The Rost invariant has trivial kernel for quasi-split groups of low rank*, Comment. Math. Helv. **76** (2001), no. 4, 684–711.
- [GH] S. Garibaldi and D.W. Hoffmann, *Totaro’s question on zero-cycles on G_2 , F_4 , and E_6 torsors*, J. London Math. Soc. (2) **73** (2006), 325–338.
- [GMS] S. Garibaldi, A.S. Merkurjev, and J.-P. Serre, *Cohomological invariants in Galois cohomology*, University Lecture Series, vol. 28, Amer. Math. Soc., 2003.
- [Gi] Ph. Gille, *Invariants cohomologiques de Rost en caractéristique positive*, K-Theory **21** (2000), 57–100.
- [HL] D.W. Hoffmann and A. Laghribi, *Quadratic forms and Pfister neighbors in characteristic 2*, Trans. Amer. Math. Soc. **356** (2004), 4019–4053.
- [J 59] N. Jacobson, *Some groups of transformations defined by Jordan algebras. I*, J. Reine Angew. Math. **201** (1959), 178–195, (= Coll. Math. Papers 63).
- [J 61] ———, *Some groups of transformations defined by Jordan algebras. III*, J. Reine Angew. Math. **207** (1961), 61–85, (= Coll. Math. Papers 65).
- [J 68] ———, *Structure and representations of Jordan algebras*, AMS Coll. Pub., vol. 39, AMS, Providence, RI, 1968.
- [J 71] ———, *Exceptional Lie algebras*, Lecture notes in pure and applied mathematics, vol. 1, Marcel-Dekker, New York, 1971.
- [J 81] ———, *Structure theory of Jordan algebras*, Lecture Notes in Mathematics, vol. 5, University of Arkansas, Fayetteville, Ark., 1981.
- [JK] N. Jacobson and J. Katz, *Generically algebraic quadratic Jordan algebras*, Scripta Math. **29** (1973), no. 3–4, 215–227.
- [KMRT] M.-A. Knus, A.S. Merkurjev, M. Rost, and J.-P. Tignol, *The book of involutions*, Colloquium Publications, vol. 44, Amer. Math. Soc., 1998.
- [Kne] M. Kneser, *Galois-Kohomologie halbeinfacher algebraischer Gruppen über p -adischen Körpern. II*, Math. Z. **89** (1965), 250–272.
- [Lam] T.Y. Lam, *Introduction to quadratic forms over fields*, Graduate Studies in Mathematics, vol. 67, American Mathematical Society, Providence, RI, 2005.
- [Lang] S. Lang, *Algebra*, third ed., Springer-Verlag, 2002.
- [Lo 75] O. Loos, *Jordan pairs*, Lecture Notes in Mathematics, vol. 460, Springer-Verlag, Berlin, 1975.
- [Lo 77] ———, *Bounded symmetric domains and Jordan pairs*, University of California-Irvine, 1977.
- [McC 66] K. McCrimmon, *A general theory of Jordan rings*, Proc. Nat. Acad. Sci. U.S.A. **56** (1966), 1072–1079.
- [McC 69] ———, *The Freudenthal-Springer-Tits constructions of exceptional Jordan algebras*, Trans. Amer. Math. Soc. **139** (1969), 495–510.
- [McC 71] ———, *Inner ideals in quadratic Jordan algebras*, Trans. Amer. Math. Soc. **159** (1971), 445–468.
- [P] H.P. Petersson, *Structure theorems for Jordan algebras of degree three over fields of arbitrary characteristic*, Comm. Alg. **32** (2004), 1019–1049.
- [PR 84] H.P. Petersson and M.L. Racine, *Springer forms and the first Tits construction of exceptional Jordan division algebras*, Manuscripta Math. **45** (1984), 249–272.
- [PR 95] ———, *On the invariants mod 2 of Albert algebras*, J. Algebra **174** (1995), no. 3, 1049–1072.
- [Ra 72] M.L. Racine, *A note on quadratic Jordan algebras of degree 3*, Trans. Amer. Math. Soc. **164** (1972), 93–103.
- [Ra 77] ———, *Point spaces in exceptional quadratic Jordan algebras*, J. Algebra **46** (1977), 22–36.
- [Sch] W. Scharlau, *Quadratic and Hermitian forms*, Springer, Berlin, 1985.

- [Spr] T.A. Springer, *Jordan algebras and algebraic groups*, Ergebnisse der Mathematik und ihrer Grenzgebiete, vol. 75, Springer-Verlag, 1973.
- [Ti 66a] J. Tits, *Algèbres alternatives, algèbres de Jordan et algèbres de Lie exceptionnelles. I. Construction*, Nederl. Akad. Wetensch. Proc. Ser. A **69** = Indag. Math. **28** (1966), 223–237.
- [Ti 66b] ———, *Classification of algebraic semisimple groups*, Algebraic Groups and Discontinuous Subgroups, Proc. Symp. Pure Math., vol. IX, AMS, 1966, pp. 32–62.
- [Ti 71] ———, *Représentations linéaires irréductibles d'un groupe réductif sur un corps quelconque*, J. Reine Angew. Math. **247** (1971), 196–220.
- [Ti 90] ———, *Strongly inner anisotropic forms of simple algebraic groups*, J. Algebra **131** (1990), 648–677.
- [V 68] F.D. Veldkamp, *Unitary groups in projective octave planes*, Compositio Math. **19** (1968), 213–258.
- [V 69] ———, *Unitary groups in Hjelmslev-Moufang planes*, Math. Z. **108** (1969), 288–312.
- [Wal] R.E. Walde, *Composition algebras and exceptional Lie algebras*, PhD thesis, University of California-Berkeley, 1967.
- [Wat] W.C. Waterhouse, *Introduction to affine groups schemes*, Graduate Texts in Mathematics, vol. 66, Springer, 1979.

DEPARTMENT OF MATHEMATICS & COMPUTER SCIENCE, EMORY UNIVERSITY, ATLANTA, GA 30322, USA

E-mail address: skip@member.ams.org

URL: <http://www.mathcs.emory.edu/~skip/>

FACHBEREICH MATHEMATIK, FERNUNIVERSITÄT IN HAGEN, D-58084 HAGEN, GERMANY

E-mail address: Holger.Petersson@FernUni-Hagen.de

URL: <http://www.fernuni-hagen.de/MATHEMATIK/ALGGE0/Petersson/petersson.html>